

DYNAMICAL PROPERTIES OF SOME ADIC SYSTEMS WITH ARBITRARY ORDERINGS

SARAH FRICK, KARL PETERSEN, AND SANDI SHIELDS

ABSTRACT. We consider arbitrary orderings of the edges entering each vertex of the (downward directed) Pascal graph. Each ordering determines an adic (Bratteli-Vershik) system, with a transformation that is defined on most of the space of infinite paths that begin at the root. We prove that for every ordering the coding of orbits according to the partition of the path space determined by the first three edges is essentially faithful, meaning that it is one-to-one on a set of paths that has full measure for every fully supported invariant probability measure. We also show that for every k the subshift that arises from coding orbits according to the first k edges is topologically weakly mixing. We give a necessary and sufficient condition for any adic system to be topologically conjugate to an odometer and use this condition to determine the probability that a random order on a fixed diagram, or a diagram constructed at random in some way, is topologically conjugate to an odometer. We also show that the closure of the union over all orderings of the subshifts arising from codings of the Pascal adic by the first edge has superpolynomial complexity, is not topologically transitive, and has no periodic points besides the two fixed points, while the intersection over all orderings consists of just four orbits.

1. INTRODUCTION

Adic, or Bratteli-Vershik (BV), systems represent all measure-preserving systems on nonatomic probability spaces and all minimal systems on the Cantor set. They provide a convenient combinatorial description of the construction of systems by the cutting and stacking method, present tail fields of stochastic processes as the fields of adic-invariant sets, facilitate the study of measure-preserving and topological orbit equivalence, and present important and interesting particular systems for detailed study. We refer to the introductions of the papers in our references list and the survey [7] for background and further details.

For adic systems, invariant measures and orbit equivalence depend only on the graph-theoretic structure of the underlying diagram and are independent of the orders assigned to the edges entering each vertex, which determine the adic transformation; further dynamical

properties, such as the spectrum, isomorphism, or joinings, do depend on the particular ordering. Some recent work has investigated how the orderings assigned to the edges can affect the sets of maximal and minimal paths. Bezuglyi, Kwiatkowski, and Yassawi [4] studied arbitrary reorderings of adic systems to determine which ones—the “perfect” orderings—allow the adic transformation to be defined as a homeomorphism. They showed that when orderings of a diagram of finite rank r are chosen uniformly at random, there is a $J \in \{1, \dots, r\}$ such that almost every ordering has J maximal and J minimal paths. Medynets [21] gave an example of a diagram for which no ordering is perfect; see [4] for generalizations. Bezuglyi and Yassawi [5] pursue this question beyond the finite rank case and give necessary and sufficient conditions for a diagram to admit a perfect ordering. Janssen, Quas, and Yassawi [18] produce a class of systems for which $J = \infty$ and almost every ordering is imperfect.

It is important to determine when coding the orbits of paths according to the natural partition into cylinder sets determined by the first k edges produces an *essentially faithful* representation of the system as a subshift on a finite alphabet; for then traditional methods of symbolic dynamics can be applied, and, even if the adic transformation cannot be defined as a homeomorphism, nevertheless the corresponding subshift can be studied as a legitimate dynamical system. By “essentially faithful” we mean that the coding map is one-to-one except on a set of paths that has measure zero for each fully supported adic-invariant probability measure on the system. This property was proved for the Pascal adic with a particular edge ordering (namely one that is lexicographical from left to right at each vertex) in [23] and for a wider class of systems in [11]. On the other hand, Downarowicz and Maass [6] (see also [17]) showed that each minimal finite topological rank adic system, namely one that can be represented up to topological conjugacy by an adic system with a bounded number of vertices on each level, with unique maximal and minimal paths, and having all orbits dense, either has a faithful coding by the first k edges for some k or else is topologically conjugate to an odometer. This result may be regarded as a generalization of previous work (see [8, 10, 16, 20]) on the relations among stationary adic systems, substitutions, and odometers. It was extended to aperiodic Cantor minimal systems in [2].

So far we have been lacking useful tests for when a general adic system allows an essentially faithful representation as a subshift or when it is topologically conjugate to an odometer. Here we show (Theorem 3.6) that for the Pascal system every edge ordering allows essentially faithful coding by the first three edges. Further, extending the ideas in [22, 23], we show that for every ordering the subshift defined by the coding is topologically weakly mixing (Theorem 4.2). We find this somewhat surprising, since the existence of eigenfunctions is highly sensitive to timing. We also establish a necessary and sufficient condition for an adic system of a fairly general kind to be topologically conjugate to an odometer (Theorem 5.3). (Previously, sufficient conditions were known for systems with a

bounded number of vertices per level [4, 8]. A similar condition has been given independently by Durand and Yassawi [9], and a necessary condition was mentioned in [13].) We use this condition to determine the probability that a random order on a fixed diagram, or an ordered diagram constructed at random in a certain sense, is topologically conjugate to an odometer (Examples 5.5 and 5.6). We also study the subshift that is the closure of the union of the subshifts determined by the Pascal adic with all possible orderings: we estimate its complexity, prove that it is not topologically transitive, and show that its only periodic points are the two fixed points. On the other hand, the intersection over all orderings consists of just four orbits. The final section presents examples and further observations related to the questions under discussion.

2. THE PASCAL SYSTEM

In this section we fix notation and terminology for the Pascal adic system. Similar concepts apply also to other adic systems.

Let P denote the Pascal graph, which is a directed graph with countably many vertices arranged into levels. The set of vertices is $\mathcal{V} = \{(x, y) : x, y \in \mathbb{Z}_{\geq 0}\}$. The set \mathcal{E} of directed edges consists of edges from (x, y) to $(x + 1, y)$ and $(x, y + 1)$ for each $(x, y) \in \mathcal{V}$. The *level* of a vertex (x, y) is $n = x + y$. For each $n \geq 0$ let

$$(1) \quad \mathcal{V}_n = \{(x, y) : x + y = n\}$$

denote the set of vertices at level n , and, for $n \geq 1$, \mathcal{E}_n denote the set of edges with source in \mathcal{V}_{n-1} and range in \mathcal{V}_n .

We specify an *ordering* of the incoming edges to all vertices other than $(0, 0)$ by a function $\xi : \mathcal{V} \rightarrow \{0, 1\}$: for $x, y > 0$, the edge from $(x - 1, y)$ to (x, y) is defined to be less than the edge from $(x, y - 1)$ to (x, y) if and only if $\xi(x, y) = 1$. As usual with adic systems, each such ordering determines a partial order on the set X of infinite directed paths on P , viewed as sequences of adjacent edges, that begin at the *root*, $(0, 0)$. Paths (γ_k) and (η_k) are *comparable* if they eventually coincide: there is an N such that $\gamma_k = \eta_k$ for all $k > N$. If n is the smallest such N , then $\gamma < \eta$ if and only if $\gamma_n < \eta_n$. Then the *adic transformation* T_ξ is defined on the complement of the set $X_{\xi, \max}$ of paths that consist entirely of maximal edges by letting $T_\xi \gamma$ equal the smallest path η that is larger than γ . Similarly, T_ξ^{-1} is defined on the complement of the set $X_{\xi, \min}$ of paths that consist entirely of minimal edges. Edges connecting “boundary vertices”, i.e. from $(x, 0)$ to $(x + 1, 0)$ and from $(0, y)$ to $(0, y + 1)$, are both maximal and minimal. Define $\gamma_{-\infty}$ to be the path down the left boundary of the diagram and γ_∞ to be the path down the right boundary. The ordinary Pascal adic system is determined by the ordering $\xi \equiv 0$. When we are working with a fixed ordering,

the subscript ξ can be omitted. Denote by X'_ξ the union of the orbits under T_ξ of all path in $X_{\xi,\min} \cup X_{\xi,\max}$. Sometimes T_ξ can be extended to a map from (part of) $X_{\xi,\max}$ to $X_{\xi,\min}$.

For each $k \geq 1$ we consider the partition \mathcal{P}_k of X into the cylinder sets

$$(2) \quad \text{Cyl}(a_1, \dots, a_k) = \{\gamma \in X : \gamma_i = a_i, i = 1, \dots, k\}$$

defined by specifying the first k edges $a_i, i = 1, \dots, k$.

These are clopen sets that generate the topology of X . For each k there are 2^k cylinder sets of level k . The elements of \mathcal{P}_k may be regarded as symbols and identified with the sets $\text{Cyl}(a_1, \dots, a_k)$, the strings $a_1 \dots a_k$, or the corresponding initial paths of length k .

The set $\mathcal{P}_k^{\mathbb{Z}}$ of two-sided infinite sequences with entries from \mathcal{P}_k is a compact metric space with distance $d(\omega, \omega') = 1/(2^{\min\{|m|: \omega_m \neq \omega'_m\}})$.

For a fixed ordering ξ , each path $\gamma \in X$ has a k -coding $\omega_k(\gamma) \in \mathcal{P}_k^{\mathbb{Z}}$ given by $\omega_k(\gamma)_n = C$ if and only if $T_\xi^n \gamma \in C \in \mathcal{P}_k$. Paths in X'_ξ have only 1-sided codings. We write $\omega \in \mathcal{P}_k^{\mathbb{Z}}$ as $\omega = \dots C_{-2}C_{-1}.C_0C_1\dots$. The subshift (Σ_k, σ) consists of the set $\Sigma_k \subset \mathcal{P}_k^{\mathbb{Z}}$ of all sequences each of whose finite subblocks can be found in $\omega_k(\gamma)$ for some $\gamma \in X$, together with the shift transformation σ defined by $(\sigma\omega)_n = \omega_{n+1}$. Σ_k is the closure of the set of all (2-sided) k -codings of all paths in $X \setminus X'_\xi$. There are natural factor maps from (Σ_k, σ) to (Σ_l, σ) for each $l < k$.

We say the k -coding is *faithful* if $\omega_k(\gamma) \neq \omega_k(\gamma')$ whenever $\gamma \neq \gamma'$. A system for which the adic transformation can be defined as a homeomorphism is expansive if and only if there is a k such that the k -coding is faithful.

The fully supported nonatomic adic-invariant ergodic probability measures on the path space X are a one-parameter family $\{\mu_\alpha : 0 < \alpha < 1\}$. Each edge from $(x, y - 1)$ to (x, y) is assigned weight α , and each edge from $(x - 1, y)$ to (x, y) is assigned weight $1 - \alpha$. The measure of a cylinder set $\text{Cyl}(a_1, \dots, a_k)$ as above is then the product of the weights on the edges that define the cylinder set. For a fixed vertex v , all cylinder sets determined by paths from $(0, 0)$ to v then have the same measure (see [23]). Changing the ordering of the edges just changes the order in which paths, or cylinder sets, are mapped to one another and does not change the set of fully supported adic-invariant ergodic probability measures.

See [3] for a result, in a completely different context, related to the following.

Proposition 2.1. *For each ordering ξ of the Pascal graph and each fully supported adic-invariant probability measure μ_α , the set $X'_\xi = \mathcal{O}(X_{\xi,\min} \cup X_{\xi,\max})$ has measure 0.*

Proof. Let $s = \max\{\alpha, 1 - \alpha\}$. For each vertex $v \in \mathcal{V}$ there is a unique (finite) reverse path from v to the root $(0, 0)$ that follows only minimal edges. (The corresponding cylinder set, which we denote by $C_{\min}(v)$, corresponds to the base of a column in the cutting and stacking representation of the system.) If v is at level n , then $\mu_\alpha(C_{\min}(v)) \leq s^n$. Thus

$$(3) \quad \mu_\alpha \left(\bigcup_{v \in \mathcal{V}_n} C_{\min}(v) \right) \leq (n+1)s^n.$$

Since each infinite path that follows only minimal edges is in the decreasing intersection of the sets $(\bigcup_{v \in \mathcal{V}_n} C_{\min}(v))$, the set of minimal paths has μ_α measure 0. Similarly, the set of maximal paths has measure 0, so X'_ξ is the countable union of sets of measure 0. \square

We assign *basic blocks* $B_\xi(x, y)$ on the alphabet $A = \{a, b\}$ to the vertices (x, y) of P as follows. The vertices $(i, 0)$, for all $i \geq 1$, are assigned the block a ; the vertices $(0, j)$, for all $j \geq 1$, are assigned the block b ; and each vertex (x, y) for which both x and y are at least 1 is assigned the block $B_\xi(x, y-1)B_\xi(x-1, y)$ if $\xi(x, y) = 0$, and $B_\xi(x-1, y)B_\xi(x, y-1)$ if $\xi(x, y) = 1$. If one of x, y is -1 , define $B_\xi(x, y)$ to be the empty block, and then this concatenation formula holds for all $x, y \geq 0$. We say that $n = x + y$ is the *level* of the basic block $B_\xi(x, y)$. The basic block $B_\xi(x, y)$ is the initial block of length $|B_\xi(x, y)|$ of the 1-coding of the minimal path from $(0, 0)$ to (x, y) . For $x, y > 0$, the number of a 's in $B_\xi(x, y)$ is given by the binomial coefficient $C(x+y-1, x-1)$, and the number of b 's by $C(x+y-1, y-1)$. If we are dealing with a fixed ordering, the subscript ξ can be omitted.

For each $k \geq 1$ basic blocks $B_\xi^k(x, y)$ on the symbols of \mathcal{P}_k of the k -coding are defined in a similar way. We retain the meaning of the coordinates x and y —the diagram is not telescoped. When $x + y = k$, for each $m = 0, 1, \dots, k$ there are $C(k, m)$ paths from the root to the vertex at $(k-m, m)$. Since they are linearly ordered by ξ , we may label them in order as $P_{k,m}^1, \dots, P_{k,m}^{C(k,m)}$. These labels $P_{k,m}^s$, for all m and s , may be taken as the 2^k symbols of the alphabet \mathcal{P}_k . For each $m = 0, 1, \dots, k$ the basic block at the vertex $(k-m, m)$ is $P_{k,m}^1 \dots P_{k,m}^{C(k,m)}$. To form the basic blocks at each level $n > k$, the level $n-1$ basic blocks are concatenated according to the ordering ξ , as above (with $B_\xi^k(x, y)$ being the empty block if one of x, y is -1). There are always two level- n basic blocks that have length 1. Note that while for $k > 1$, the corresponding alphabet for Σ_k is not $\{a, b\}$, we can still talk about counts of a 's and b 's associated to $B_\xi^k(x, y)$ by considering the factor map $\Sigma_k \rightarrow \Sigma_1$. Not only does each basic block $B_\xi^k(x, y)$ determine the vertex (x, y) , but also the k -symbol counts in $B_\xi^k(x, y)$ determine (x, y) . Note that for all x, y $B_\xi^1(x, y) = B_\xi(x, y)$ as defined in the preceding paragraph.

3. THE PASCAL SYSTEM WITH ANY ORDERING IS ESSENTIALLY EXPANSIVE

For any k and any path γ in an arbitrary adic system, due to the structure of the diagram the k -coding of the orbit of γ has a natural hierarchical structure of factorings for each $n \geq k$ into level- n basic blocks $B_\xi^k(x, y)$ with $x + y = n$. This is evident for substitution subshifts and is especially simple (periodic) for odometers. When an iterate of γ under T enters the cylinder determined by a minimal path from the root to a vertex v at level n , the k -coding at that instant begins an occurrence of a level- n basic block. (In the cutting and stacking version, the orbit of γ enters the base of a column and proceeds up the column.) In the following we continue to focus on the Pascal graph.

Fix an ordering ξ of the Pascal graph P , a coding length $k \geq 1$, and a sequence $\omega_k \in \mathcal{P}_k^\mathbb{Z}$. For each $n \geq k$ the basic blocks of level n are distinct (since, after applying the factor map $\Sigma_k \rightarrow \Sigma_1$ in case $k > 1$, each block has a different count of a and b as described in the previous section), and thus for fixed $n = x + y$, $B_\xi^k(x, y)$ determines the vertex (x, y) of P to which it is associated. We are interested in hierarchical factorings of sequences $\omega_k \in \Sigma_k$ into concatenations of basic blocks so that the factorizations are consistent across levels, respect the ordering ξ , and are nontrivial. This idea is made precise in the following definition.

Definition 3.1. Given $\omega_k \in \Sigma_k$, a ξ -consistent factoring scheme for ω_k (more briefly, a factoring scheme, or scheme) is a sequence of factorings of ω_k into basic blocks of level n for each $n \geq k$ as follows. For each $n \geq k$, ω_k is written as a concatenation $\dots W_{n,-1}W_{n,0}W_{n,1}W_{n,2}\dots$ of blocks $W_{n,i}$, in such a way that

- (1) each $W_{n,i}$ is a basic block $B_\xi^k(x_{n,i}, y_{n,i})$ of level n with respect to ξ ;
- (2) there is a function $j : \mathbb{N} \times \mathbb{Z} \rightarrow \mathbb{Z}$ such that $j(n, i + 1) = j(n, i) + 2$ for all $n \geq k$ and all i , and, for each $n > k$ and each i , $W_{n,i} = W_{n-1, j(n,i)}W_{n-1, j(n,i)+1}$;
- (3) if $\xi(x_{n,i}, y_{n,i}) = 0$, then $W_{n-1, j(n,i)} = B_\xi^k(x_{n,i}, y_{n,i} - 1)$ and $W_{n-1, j(n,i)+1} = B_\xi^k(x_{n,i} - 1, y_{n,i})$, while if $\xi(x_{n,i}, y_{n,i}) = 1$, then $W_{n-1, j(n,i)} = B_\xi^k(x_{n,i} - 1, y_{n,i})$ and $W_{n-1, j(n,i)+1} = B_\xi^k(x_{n,i}, y_{n,i} - 1)$;
- (4) there is no interval $[i, i + j] \subset \mathbb{Z}$ with $j \geq 1$ such that each $W_{n,r}$, for $i \leq r \leq i + j$, has length 1 and the string $W_{n,i} \dots W_{n, i+j}$ is a level- n basic block;
- (5) denoting for each $n \geq k$ by $CB(n) = (\omega_k)_{-i_n} \dots (\omega_k)_{j_n}$ the level- n basic block of the factorization that includes the coordinate 0, we have $i_n, j_n \rightarrow \infty$ for all i .

Condition (4) is needed to ensure that no part of ω_k is factored trivially into length 1 level- n basic blocks unless necessary. Condition (5) is needed to avoid sequences such as those in the remark below. Note that for every k the k -coding of every path in X (even if one-sided, since we allow empty basic blocks when a coordinate is negative) has a natural

factoring scheme determined by the sequence of vertices through which it passes, but one-sided sequences do not satisfy (5). In principle, for some k a k -coding of a path might have more than one consistent factoring scheme. In the following Proposition we show that each sequence with a consistent factoring scheme determines a unique corresponding path. In Theorem 3.6 and Corollary 3.7 we show that the 3-coding of every path in $X \setminus X'_\xi$ has a unique consistent factorization scheme since the 3-coding corresponds to a unique path.

Remark 3.2. (1) The subshift (Σ_k, σ) consists of the *closure* of the (therefore dense) set of k -codings of paths in $X \setminus X'_\xi$ and thus may contain sequences ω_k that are not the codings of any paths. For example, in the Pascal system with the usual ordering $\xi \equiv 0$, for every n the block $ab^n a^n b$ appears in the 1-coding of some path in $X \setminus X'_\xi$ (these blocks are found inside basic blocks near the centers of rows of the diagram), and hence the sequence $\omega = b^\infty . a^\infty$ is in Σ_1 . If for this sequence we allowed the (unique) trivial factorization into blocks of length 1 to count as a consistent factoring scheme, in order to find a path with this coding (guaranteed by Proposition 3.3) we would try to define T_ξ on γ_∞ and $\gamma_{-\infty}$, where it was not defined before, in such a way that ω would become the 1-coding of the orbit of γ_∞ . But trying to define $T_\xi \gamma_\infty = \gamma_{-\infty}$, $T_\xi \gamma_{-\infty} = \gamma_{-\infty}$ (the only possibilities) fails, because then we would need $T_\xi^{-1} \gamma_\infty$ equal to both γ_∞ and $\gamma_{-\infty}$ and $T_\xi^{-1} \gamma_{-\infty}$ equal to both γ_∞ and $\gamma_{-\infty}$.

(2) At this time we do not know whether there could be a (two-sided) sequence in some Σ_k that (a) is not the k -coding of a path in $X \setminus X'_\xi$; (b) has a unique factoring scheme satisfying (1)–(4) above plus a weaker version of (5), in which it is required only that at least one of i_n, j_n tends to infinity; and (c) allows T_ξ to be defined on some maximal and minimal paths in such a way that the sequence becomes the coding of the new orbit. Theorem 3.6 can be generalized to cover this possibility. Working with the usual ordering $\xi \equiv 0$, the sequence $a^\infty . ba^\infty \in \Sigma_1$ satisfies (a) and (b), but not (c). The path γ that passes through $(0,0)(1,0)(1,1)(2,1)(3,1)(4,1) \dots$ has its backward orbit code to a^∞ (every entry to the left of the center is a), but then $T_\xi \gamma$ is maximal, $T_\xi^2 \gamma$ must be defined to be the left edge $\gamma_{-\infty}$, we have to define $T_\xi \gamma_{-\infty} = \gamma_{-\infty}$, and T_ξ^{-1} cannot be well defined.

Proposition 3.3. *Suppose that an ordering ξ of the Pascal graph P , $k \geq 1$, and a sequence $\omega \in \mathcal{P}_k^\mathbb{Z}$ with a ξ -consistent factoring scheme are given. Then there is a unique path $\gamma \in X \setminus X'_\xi$ such that the k -coding of the orbit of γ under the adic transformation T_ξ is the given sequence ω and the natural factoring defined by the path coincides with the given one.*

Proof. Suppose that an ordering ξ and $\omega \in \mathcal{P}_k^\mathbb{Z}$ with a ξ -consistent factoring scheme are given. Denote in the following by CB_n the central basic block of level n in the given factorization, that is, the one that includes the coordinate 0. Let $k_n \in \{0, 1, \dots, |CB_n| - 1\}$ be such that $\sigma^{-k_n} \omega$ has origin at the beginning of CB_n . Form a path γ which at each level n is at vertex v_n determined by the level- n central block CB_n , and with initial segment of

length n equal to $T_\xi^{k_n}$ applied to the minimal path from $(0,0)$ to v_n . The central symbol of ω determines the vertices of γ at levels $0, \dots, k$ for $n \geq k$. Then the k -coding $\omega_k(\gamma)$ of γ equals the given sequence ω on the range of coordinates $I_n = [-k_n, |CB_n| - 1 - k_n]$ for every n . Thus the coding and factorization determine the sequence of vertices through which the path passes, and no other path can have the same coding and natural factorization. Since $\cup\{I_n : n = 1, 2, \dots\} = \mathbb{Z}$, we have $\gamma \in X \setminus X'_\xi$ and the full coding $\omega_k(\gamma)$ equals the full sequence ω . \square

Lemma 3.4. *Fix an ordering ξ of the Pascal diagram, with path space X . Suppose γ, γ' are distinct paths in $X \setminus X'_\xi$. Suppose further that γ and γ' follow only minimal edges from $(0,0)$ to distinct vertices (x,y) and (x',y') respectively, where $x+y = x'+y' = n$ for some $n \geq 1$. Then the 3-codings of γ and γ' are not the same. In other words, two such paths that follow only minimal edges starting at the root to different vertices at the same level do not have the same coding by initial path segments of length 3.*

Proof. Suppose γ and γ' are as above. In particular, we may assume that the 1- (and 2-) codings of γ and γ' are the same: $\omega_1(\gamma) = \omega_1(\gamma')$ and $\omega_2(\gamma) = \omega_2(\gamma')$ (otherwise we are done). The longest minimal path from $(0,0)$ that is contained in both γ and γ' ends at some vertex $v = (x_0, y_0)$ at level $n_0 = x_0 + y_0$. In other words, γ and γ' *split* at v . If $n_0 < 3$ then γ and γ' would disagree on one of their first three edges and we would be done; we therefore consider only the cases where $n_0 \geq 3$. We may assume that no paths with a lower level splitting occur (at the same time) in the forward orbits of γ and γ' , respectively. Specifically, find a smallest n_0 for which there exists an r such that $T^r\gamma$ and $T^r\gamma'$ are minimal until distinct vertices at level $n_0 + 1$ and contain the same edges until level n_0 , replace γ and γ' by $T^r\gamma$ and $T^r\gamma'$, and define v to be the vertex where this latter pair split. It is the minimality of n_0 that we will exploit in the following proof by contradiction.

We consider several cases according to where in the diagram the vertex v can occur. Our first case is when v is an interior vertex. In other words, let $v \in U = \{(x,y) : x > 0 \text{ and } y > 0\}$. Since both paths are minimal until level $n > n_0$, there exist two minimal edges with source v ; label γ, γ' so that the left one is contained in γ and the right one contained in γ' . Let k_1 be the length of the basic block at v and consider the paths $T^{k_1}\gamma$ and $T^{k_1}\gamma'$. Then at level n_0 , $T^{k_1}\gamma$ passes through vertex $(x_0 + 1, y_0 - 1)$ and $T^{k_1}\gamma'$ passes through vertex $(x_0 - 1, y_0 + 1)$, and both are minimal until level $n_0 = x_0 + y_0$. Therefore the split between the two paths in the orbits of γ and γ' occurred at some level n_1 where $n_1 \leq n_0 - 2$. This contradicts the minimality of n_0 . (See Figure 1, where we indicate minimal edges by dashed lines and maximal edges by solid lines.)

Now consider the case when v is on the left side of the diagram (i.e. $v = (x_0, 0)$ for some $x_0 \geq 3$). Since $\gamma, \gamma' \notin \{\gamma_{-\infty}, \gamma_\infty\}$ ($\gamma_{-\infty}$ and γ_∞ do not have a ξ -consistent coding scheme),




FIGURE 1. $T^{k_1}\gamma$ and $T^{k_1}\gamma'$ do not agree into level $n_0 = x_0 + y_0$.

both γ and γ' must leave the left side of the diagram. Consider first the case when both leave the left side via minimal edges. See Figure 2. Then $n_0 = x_0$ and we label so that γ contains the minimal path to $(x_0 + 1, 0)$ and γ' contains the minimal path to $(x_0, 1)$. Since γ leaves the left side of the diagram, it passes through a vertex $(x_1, 1)$, and this edge is minimal. In this case, we will examine the first edge coding of γ and γ' . We first note that $B(x_0, 1) = a^{i_0}ba^{j_0}$ where $i_0 + j_0 = x_0$. (In general, the basic block at $(x, 1)$ for any $x > 0$ is of the form $a^i ba^j$ for some $i + j = x$, and i and j are nondecreasing as x increases.) Since the edge connecting vertices $(x_1, 0)$ and $(x_1, 1)$ is minimal, $B(x_1, 1) = aB(x_1 - 1, 1)$. Since $x_0 < x_1$, $B(x_1 - 1, 1)$ contains $B(x_0, 1)$ (the blocks are the same if $x_1 = x_0 + 1$). Therefore $B(x_1, 1) = a^{i_1}ba^{j_1}$ with $i_1 > i_0$ and $j_1 \geq j_0$. Since γ and γ' are minimal into $(x_1, 1)$ and $(x_0, 1)$ respectively, $a^{i_1}ba^{j_1}$ immediately follows the decimal in $\omega_1(\gamma)$ (i.e. beginning in the origin position) and $a^{i_0}ba^{j_0}$ immediately follows the decimal in $\omega_1(\gamma')$. However, $i_1 > i_0$ contradicts our assumption that γ and γ' have the same 1-coding.




FIGURE 2. The dotted lines are necessarily minimal and the thick lines are necessarily maximal.

(An analogous argument rules out the case where γ and γ' split along the diagonal $x = 0$ and contain minimal paths from the root to $x = 1$.)

Our final and most complex case is the case when v is on the left side of the diagram and γ' leaves along a minimal edge while γ leaves along a maximal edge at some later level. See Figure 3. Thus $v = (x_0, 0)$, for some $x_0 \geq 3$, $n_0 = x_0$, γ contains the minimal path to $(x_0 + 1, 0)$, γ' contains the minimal path to $(x_0, 1)$ and γ exits the left side at $(x_1, 0)$, for some $x_1 > x_0$. If γ is not in $X_{\xi, \max}$ it does not continue along maximal edges forever. Let (u, w) at level $m = u + w$ be the range of the first minimal edge in γ after $(x_1, 1)$. Note that $m \geq x_1 + 2 \geq x_0 + 3$. Since the edge connecting $(x_0, 0)$ and $(x_0, 1)$ is the first non-maximal edge in γ' , $T\gamma'$ passes through vertex $(x_0 - 1, 1)$. In particular, $T\gamma'$ contains the minimal path from the root vertex into $(x_0 - 1, 1)$. In order not to contradict the minimality of n_0 , $T\gamma$ must also pass through vertex $(x_0 - 1, 1)$. Since γ did not pass through $(x_0, 1)$, $T\gamma$ must contain the edge connecting $(x_0 - 1, 1)$ and $(x_0 - 1, 2)$ and this edge must be minimal according to the edge ordering. See Figure 3. Thus $T\gamma$ actually consists of all minimal edges up to level $m - 1 \geq x_0 + 2$ and therefore either passes through $(x_0 - 1, 3)$ or $(x_0, 2)$ along a minimal edge. It is this fact that we will use to arrive at a contradiction.




FIGURE 3. The dotted lines are necessarily minimal and the thick red lines are necessarily maximal.

We now consider $T^{k_1}\gamma$ where $k_1 = |B(x_0 - 1, 1)| + 1$, in other words, the path in the forward orbit of $T\gamma$ which first contains the maximal edge from $(x_0 - 2, 2)$ to $(x_0 - 1, 2)$. It contains the unique minimal path from $(0, 0)$ to $(x_0 - 2, 2)$. By an argument similar to the above, we now know that $T^{k_1}\gamma'$ must also contain the unique minimal path from $(0, 0)$ to

$(x_0 - 2, 2)$ (since otherwise the orbits of γ and γ' would split earlier than level n_0), but then continue along the edge with range $(x_0 - 2, 3)$, which is necessarily minimal. See Figure 4.




FIGURE 4. Step 2.

The issue comes at the next stage when we consider $T^{k_2}\gamma$ and $T^{k_2}\gamma'$, where $k_2 = k_1 + |B(x_0 - 2, 2)|$. We see that $T^{k_2}\gamma'$ passes through vertex $(x_0 - 3, 3)$. However, since the edge contained in $T^{k_1}\gamma$ leaving vertex $(x_0 - 1, 2)$ is minimal, we see from the ordering on the edges that has been established thus far that $T^{k_2}\gamma$ must pass either through vertex $(x_0, 0)$ or vertex $(x_0 - 2, 2)$ at level $n_0 = x_0$. See Figure 5. Therefore we again contradict the minimality of n_0 .




FIGURE 5. The two possibilities for $T^{k_2}\gamma$

□

Remark 3.5. Replacing T by T^{-1} in the above argument shows that two paths that are maximal until different vertices at the same level also do not have the same coding by initial segments of length 3.

Theorem 3.6. *Fix an ordering ξ of the Pascal diagram, with path space X . Suppose γ, γ' are distinct paths in $X \setminus X'_\xi$. Then the 3-codings of γ and γ' are not the same. In other words, for such paths the 3-coding is faithful.*

Proof. Suppose, to the contrary, that there are two such distinct paths γ and γ' whose codings by initial segments of length 3 are the same. There exists an $r \geq 0$ such that $T^{-r}\gamma$ and $T^{-r}\gamma'$ share the same minimal path until vertex $v = (x_0, y_0)$, for some x_0, y_0 with $n_0 = x_0 + y_0 \geq 3$, and then follow different edges out of that vertex. Replacing γ and γ' by $T^{-r}\gamma$ and $T^{-r}\gamma'$, we may assume that γ and γ' have this property. If v is on the left or right side of the diagram, then γ and γ' are minimal (or maximal) to different vertices, and so, by Lemma 3.4 (or Remark 3.5 respectively), this cannot happen. Thus $x_0, y_0 > 0$; in other words, v is an “internal” vertex.

If the first edges after v in γ and γ' are both maximal, then either $T^{-1}\gamma$ contains a maximal path from $(0, 0)$ to $(x_0 + 1, y_0 - 1)$ and $T^{-1}\gamma'$ contains a maximal path to $(x_0 - 1, y_0 + 1)$, or vice versa. It then follows from Remark 3.5 that the 3-codes for $T^{-1}\gamma$ and $T^{-1}\gamma'$ (and hence for γ and γ') cannot be the same. If the first edges after v in γ and γ' are both minimal, then by Lemma 3.4 γ and γ' have different 3-codings.

So, without loss of generality, we may assume that the first edge in γ after v is maximal and the first in γ' is minimal. Assume further that the coordinate axes are chosen so that γ goes from v to $(x_0 + 1, y_0)$ and γ' goes from v to $(x_0, y_0 + 1)$.

Now $T^{|B(x_0, y_0)|}\gamma'$ begins with a minimal path from $(0, 0)$ to $(x_0 - 1, y_0 + 1)$ (see Figure 6), so $T^{|B(x_0, y_0)|}\gamma$ has the same property. (If these paths were at different vertices at level n_0 , Lemma 3.4 would tell us that they had distinct 3-codings.) So these paths pass through the vertex $(x_0 - 1, y_0 + 1)$. Note that if γ is not in the orbit of a maximal path, then there is a finite string of maximal edges in γ descending from v and followed by a single minimal edge from a vertex $v_1 = (x_1, y_1)$ to a vertex $v_2 = (x_2, y_2)$.

We claim that the next edge in γ' after $(x_0, y_0 + 1)$ is minimal. If this were not the case, then $T^{-1}\gamma'$ would contain a maximal path from $(0, 0)$ to either $(x_0 - 2, y_0 + 2)$ or (x_0, y_0) , whereas $T^{-1}\gamma$ would contain a maximal path to $(x_0 + 1, y_0 - 1)$. See Figure 7. Again, this would mean that the 3-codings for γ and γ' are different.

So the next edge in γ' after $(x_0, y_0 + 1)$ is minimal. Letting

$$(4) \quad r = |B(x_0, y_0)| + |B(x_0 - 1, y_0 + 1)|,$$




FIGURE 6

it follows that $T^r\gamma$ contains a minimal path from $(0, 0)$ to $(x_0 - 2, y_0 + 2)$, whereas $T^r\gamma'$ contains a minimal path to either $(x_0 + 1, y_0 - 1)$ or $(x_0 - 1, y_0 + 1)$. See Figure 8. This is ruled out by applying the Lemma to $T^r\gamma$ and $T^r\gamma'$. \square

Corollary 3.7. *For each ordering ξ of the Pascal graph P and each path $\gamma \in X \setminus X'_\xi$, the 3-coding $\omega_3(\gamma) \in \mathcal{P}_k^{\mathbb{Z}}$ has a unique ξ -consistent factoring scheme.*

Proof. Fix a particular ordering ξ on the Pascal graph P . That for each $\gamma \in X \setminus X'_\xi$, $\omega_3(\gamma)$ has such a natural factoring scheme is clear from the definitions of $\omega_3(\gamma)$ and the basic blocks $B_\xi^3(x, y)$, as mentioned previously. By Proposition 3.3 there is a unique path that has the same 3-coding and natural factoring scheme. Another factoring of the 3-coding of γ would determine a different path which would have the same 3-coding as γ , contradicting Theorem 3.6. \square

Remark 3.8. The preceding Corollary helps to clarify the relationship among paths, factorizations of codings, and factorizations of basic blocks. Suppose we have an ordering ξ on any Bratteli diagram and know that there is a subset X'_ξ of the path space X such that for each path $\gamma \in X \setminus X'_\xi$ the k -coding $\omega_k(\gamma)$ admits a unique factorization scheme into basic blocks. (Perhaps X'_ξ has measure 0 for each fully supported ergodic invariant measure.) Suppose further that the basic blocks at each level are distinct. Then it follows that distinct




FIGURE 7

paths in $X \setminus X'_\xi$ have distinct k -codings, since, as in the argument above, the factorization scheme would determine the path. Further, each basic block $B_\xi^k(x, y)$ would have a factorization into basic blocks of level n for each n with $k \leq n < x + y$, unique if without trivial segments as in the definition of factorization scheme. Basic blocks of level $l < k$ might not have unique factorizations; the central symbol specifies the path up to the root and hence a preferred factorization of the basic blocks of levels $l < k$. Thus under such hypotheses we would have unique factorization of basic blocks of levels greater than or equal to k , as shown for the case of the Pascal diagram, 1-codings, and special orderings in Lemma 6.1.

4. TOPOLOGICAL WEAK MIXING

Recall that for a fixed ordering ξ , (Σ_k, σ) denotes the subshift on 2^k symbols generated by coding paths according to their first k edges. This is a legitimate topological dynamical system—a compact metric space together with a homeomorphism. (We use the metric $d(\omega, \omega') = 1/2^{\inf\{|n|:\omega(n) \neq \omega'(n)\}}$). By Theorem 3.6 the coding is essentially faithful if $k \geq 3$. Denote by $\mathcal{B}(\Sigma_k)$ the set of bounded complex-valued functions on Σ_k whose set of continuity




FIGURE 8

points is residual. In order to prove that (Σ_k, σ) is topologically weakly mixing, we follow the plan in [22, 23] to check the condition of Keynes and Robertson [19]: a topological dynamical system which is “closed ergodic” (there is an invariant Borel probability measure on the space for which every closed invariant set has measure 0 or 1) is topologically weakly mixing if and only if every $f \in \mathcal{B}(\Sigma_k)$ for which there is $\lambda \in \mathbb{C}$ such that $f \circ \sigma = \lambda f$ everywhere must be constant on a dense set. (It is not enough to show that there are no nonconstant *continuous* eigenfunctions, since the subshift is not minimal.) For this purpose we have to go beyond the polynomial equidistribution theorem of Weyl and quote instead a theorem of F. Hahn [14].

In our situation (Σ_k, σ) is closed ergodic, since there are fully supported ergodic measures on Σ_k (the measures $\mu_\alpha, 0 < \alpha < 1$, mentioned above). Our system is topologically ergodic (there exist dense orbits). We aim to prove that if $f \in \mathcal{B}(\Sigma_k)$ and $\lambda \in \mathbb{C}$ satisfy $f \circ \sigma = \lambda f$ everywhere, then $\lambda = 1$. Then it will follow from another theorem of Keynes and Robertson [19] (since then $f \circ \sigma = f$ everywhere) that f is constant on a dense set, and consequently the system is topologically weakly mixing. We also use the fact [23, Lemma 4.18] that if $f \in \mathcal{B}(\Sigma_k)$ and $\lambda \in \mathbb{C}$ satisfy $f \circ \sigma = \lambda f$ everywhere, then every point which has both its

forward and backward orbit dense is a point of continuity of f . Another essential ingredient is the following extension of the “Kink Lemma” of [22, 23] taking into account the unknown ordering of edges.

Lemma 4.1. *Let X denote, as above, the set of infinite directed paths on the Pascal graph, and fix an ordering ξ . Let $\gamma = (\gamma_r) \in X$, with γ_r denoting the edge of γ from level $r - 1$ to level r , be a path to a vertex (i, j) , with $i > 0$ and $j > 0$, which continues to the vertex $(i + 1, j + 1)$ in such a way that the edge γ_{i+j+2} of γ that enters $(i + 1, j + 1)$ is minimal. Let $n = i + j \geq k$ and denote by e the edge with source (i, j) that is not equal to γ_{n+1} . We have to consider eight cases, labeled by triples (a_1, a_2, a_3) , depending on whether γ_{n+1} is maximal ($a_1 = \max$) or minimal ($a_1 = \min$), e is maximal ($a_2 = \max$) or minimal ($a_2 = \min$), and γ_{n+1} has range $(i + 1, j)$ ($a_3 = LR$) or $(i, j + 1)$ ($a_3 = RL$). Let*

$$(5) \quad r_n = \begin{cases} C(n, j) & \text{cases } (\max, \min, LR) \text{ and } (\max, \min, RL) \\ C(n + 1, j + 1) & \text{cases } (\min, \min, RL) \text{ and } (\max, \max, LR) \\ C(n + 1, j) & \text{cases } (\min, \min, LR) \text{ and } (\max, \max, RL) \\ C(n + 1, j) + C(n, j + 1) & \text{cases } (\min, \max, LR) \text{ and } (\min, \max, RL). \end{cases}$$

Then $T^{r_n}\gamma$ and γ agree on their first n edges. Consequently, if x is a path that begins with the minimal path to (i, j) , z is a path that begins with the maximal path to (i, j) , $T^m x = \gamma$, and $T^l \gamma = z$, then $d(\sigma^{r_n} \omega_k(\gamma), \omega_k(\gamma)) \leq 1/2^{\min\{m, l\}}$.

Proof. Let z denote the path which is maximal to the vertex (i, j) and agrees with γ after level $n = i + j$, so that $z = T^l \gamma$ for some $l \geq 0$. Suppose first that the edge γ_{n+1} is maximal and e is minimal. Then in both cases (\max, \min, LR) and (\max, \min, RL) (whether γ_{n+1} has range $(i + 1, j)$ or $(i, j + 1)$) Tz begins with the minimal path to (i, j) . Then choosing $m \geq 0$ so that $m + l + 1 = C(n, j)$ yields that $T^m Tz = T^{m+l+1} \gamma$ agrees with γ on their first n edges. Then $r_n = C(n, j)$ has the stated property. See Figure 9.

The other cases are similar but differ in the details, and the particular values of r_n that we find are important for the following theorem, so we go through them all. Suppose now that the edge e is maximal and γ_{n+1} is maximal and has range $(i + 1, j)$ (case (\max, \max, LR)). In this case Tz begins with the minimal path to the vertex $(i - 1, j + 1)$ followed by the minimal edge to $(i, j + 1)$ and the maximal edge to $(i + 1, j + 1)$. Then $u = T^{C(n, j+1)-1} Tz$ begins with the maximal path to $(i - 1, j + 1)$, and $y = Tu$ begins with the minimal path to (i, j) followed by the maximal edge to $(i, j + 1)$. Choose $m \geq 0$ so that $m + l + 1 = C(n, j)$. Then $T^m Tu$ agrees with γ on their first n edges. We have $T^m Tu = T^m T^{C(n, j+1)-1} Tz = T^{m+l+1+C(n, j+1)} \gamma = T^{C(n, j)+C(n, j+1)} \gamma = T^{C(n+1, j+1)} \gamma$, so $r_n = C(n + 1, j + 1)$ acts as stated.




FIGURE 9. Case (max, min, LR)

If the edge e is maximal and γ_{n+1} is maximal and has range $(i, j+1)$ (case (max, max, RL)), the argument is the same as the immediately preceding one, except that j is replaced by $j-1$. Namely, Tz begins with the minimal path to $(i+1, j-1)$, $u = T^{C(n, j-1)-1}Tz$ begins with the maximal path to $(i+1, j-1)$, $y = Tu$ begins with the minimal path to (i, j) followed by the maximal edge to $(i+1, j)$. Again choose $m \geq 0$ so that $m+l+1 = C(n, j)$. Then T^mTu and γ agree on their first n edges, so we obtain $r_n = C(n, j) + C(n, j-1) = C(n+1, j)$.

Consider now the case (min, max, LR). Then Tz begins with the minimal path to $(i+1, j-1)$, $u = T^{C(n, j-1)-1}Tz$ begins with the maximal path to $(i+1, j-1)$, and Tu begins with the minimal path to $(i-1, j+1)$ followed by a minimal and then a maximal edge. Then $y = TT^{C(n, j+1)-1}Tu$ begins with the minimal path to (i, j) extended by two maximal edges. Thus choosing $m \geq 0$ so that $m+l+1 = C(n, j)$ yields that $T^my = T^{m+l+1+C(n, j+1)+C(n, j-1)}\gamma = T^{C(n+1, j)+C(n, j+1)}\gamma$ agrees with γ on the first n edges.

Next, consider the case (min, max, RL). Then Tz is minimal to $(i-1, j+1)$, $u = T^{C(n, j+1)-1}Tz$ is maximal to $(i-1, j+1)$, and $p = Tu$ is minimal to $(i+1, j)$ extended by a maximal edge. Then $T^{C(n, j-1)-1}p$ is maximal to $(i+1, j-1)$, so that $y = T^{C(n, j-1)}p$ is minimal to (i, j) . Thus if we choose m so that $m+l+1 = C(n, j)$, we will have $T^my = T^mT^{C(n, j-1)}p = T^{m+C(n, j-1)+C(n, j+1)+l+1}\gamma = T^{C(n+1, j)+C(n, j+1)}\gamma$ and γ agreeing on their first n edges.

Now consider the case (min, min, LR). Then Tz starts with the minimal path to $(i+1, j-1)$, $u = T^{C(n, j-1)-1}Tz$ starts with the maximal path to $(i+1, j-1)$, $y = Tu$ is

minimal to (i, j) followed by a minimal and then a maximal path; and if we choose m so that $m + l + 1 = C(n, j)$, then $T^m y = T^{C(n+1, j)} \gamma$ and γ agree on the first n edges.

The final case is (min, min, RL). Then Tz is minimal to $(i - 1, j + 1)$ and extends from there by a maximal edge followed by a minimal edge. Then $u = T^{C(n, j+1)-1} Tz$ is maximal to $(i - 1, j + 1)$, from where it extends with a maximal and then a minimal edge. Thus $y = Tu$ is minimal to (i, j) and extends by the minimal edge e followed by a maximal edge; and, if we choose m so that $m + l + 1 = C(n, j)$, then $T^m y = T^{m+l+1+C(n, j+1)} = T^{C(n+1, j+1)} \gamma$ and γ agree on their first n edges.

The final statement in the Lemma follows from the fact that $T^i(T^{r_n} \gamma)$ and $T^i \gamma$ agree on their first $n \geq k$ edges for $-m \leq i \leq l$. \square

Theorem 4.2. *For each ordering, the subshift (Σ_k, σ) is topologically weakly mixing.*

Proof. Fix an ordering ξ . We deal only with paths in X whose orbit does not contain any paths that are eventually diagonal or eventually consist of only maximal or only minimal edges. (A path with vertices (x_k, y_k) , $k \geq 0$, is *eventually diagonal* if either coordinate is eventually constant. The set of such paths is countable.) By Proposition 2.1 the set of such paths has measure 1 for each fully supported ergodic T -invariant probability measure on X . The coding $\omega_k(\gamma)$ of any such path γ has both its forward and backward orbit dense in Σ_k and hence, as noted above, is a point of continuity of any eigenfunction $f \in \mathcal{B}(\Sigma_k)$.

Suppose that $f \in \mathcal{B}(\Sigma_k)$, $\lambda \in \mathbb{C}$, and $f \circ \sigma = \lambda f$ everywhere on Σ_k ; we want to show that then $\lambda = 1$. Fix a point $\omega_k(\gamma)$ that is the coding of a path γ as described above and hence is a point of continuity of f . For each n let (i_n, j_n) denote the vertex through which γ passes at level n , i.e. the range of the edge γ_n . For each n , the path γ can be modified if necessary below level n to a path γ' that satisfies the hypotheses of Lemma 4.1, is not eventually diagonal, and does not eventually consist of only maximal or minimal edges. This determines r_n, m_n, l_n so that $T^{r_n} \gamma', \gamma'$, and γ agree on their first n edges, and $d(\sigma^{r_n} \omega_k(\gamma'), \omega_k(\gamma')) \leq 1/2^{\min\{m_n, l_n\}}$. Since m_n and l_n are unbounded this implies that $d(\sigma^{r_n} \omega_k(\gamma'), \omega_k(\gamma')) \rightarrow 0$ as $n \rightarrow \infty$. Since $f(\sigma^{r_n} \omega_k(\gamma')) = \lambda^{r_n} f(\omega_k(\gamma'))$ for all n , we have $\lambda^{r_n} \rightarrow 1$ as $n \rightarrow \infty$. In particular $|\lambda| = 1$.

Suppose that λ is a root of unity, so that $\lambda = \exp(2\pi ia/b)$ for two relatively prime integers a and b . Let q be a prime that divides b . In the Pascal triangle we can find infinitely many n for which there is a j such that none of $C(n, j), C(n+1, j+1), C(n+1, j), C(n+1, j) + C(n, j+1)$ is divisible by q . For example, by Lemma 5.1 of [1] $C(q^s - 2, k) \equiv_q (-1)^k (k+1)$ for $k = 0 \dots, q^s - 2$. Therefore, for each s the entire row $C(q^s - 1, j), 0 \leq j \leq q^s - 1$, is not divisible by q , and in the preceding row there are many entries $C(q^s - 2, tq - 1), 0 < t \leq q^{s-2}$,

that are not divisible by q as well as many that are, so choosing $n = q^s - 2$ and j so that $j + 1$ is not divisible by q but $j + 2$ is divisible by q will do. (If $j + 1$ is not divisible by q , then neither is $C(q^s - 2, j)$, while if $j + 2$ is divisible by q , then so is $C(q^s - 2, j + 1)$.) We can arrange that our path γ hits such vertices for infinitely many n , and then $|\lambda^{r_n} - 1| \geq 1/q$ for all of these values of n , contradicting that $\lambda^{r_n} \rightarrow 1$. Thus there are no roots of unity besides 1 among the eigenvalues of σ .

Suppose now that λ is not a root of unity, say $\lambda = \exp(2\pi i\beta)$ for an irrational $\beta \in (0, 1)$. According to the theorem of F. Hahn [14, Theorem 5] the transformation $S(x_1, x_2, x_3, \dots) = (x_1 + \beta, x_2 + x_1, x_3 + x_2, \dots) \pmod{1}$ on the infinite torus is minimal, in fact strictly ergodic, with Haar measure the unique invariant probability measure. Then the orbit of $(0, 0, 0, \dots)$ is dense, and $S^n(0, 0, 0, \dots) = (C(n, 1)\beta, C(n, 2)\beta, \dots, C(n, n)\beta, 0, 0, \dots) \pmod{1}$. We use this result to construct a continuity point $\omega_k(\gamma)$ of f that is the coding of a path γ such that, denoting by (i_n, j_n) the vertex through which γ passes at level n , there are infinitely many n for which all of $C(n, j_n)\beta \pmod{1}, C(n + 1, j_n + 1)\beta \pmod{1}, C(n + 1, j_n)\beta \pmod{1}$, and $[C(n + 1, j) + C(n, j + 1)]\beta \pmod{1}$ are a fixed distance from 0 and 1, again making it impossible that $\lambda^{r_n} \rightarrow 1$. Given an initial segment of γ ending at a vertex (i_0, j_0) , extend γ several steps so as to use both some maximal and some minimal edges as well as some edges that increase the first (i) coordinate and the second (j) coordinate, arriving at a vertex (i_1, j_1) . Then add edges that increase the i coordinate until we arrive at a vertex (i, j) for which $(C(i + j, j - 1), C(i + j, j), C(i + j, j + 1))\beta \approx (1/8, 1/8, 1/8) \pmod{1}$ (say $\{|C(i + j, j - 1)\beta - 1/8| < 1/16, \text{ etc.}, \text{ with } \{x\} \text{ denoting } x \pmod{1}\}$). Then $C(i + j + 1, j)\beta \pmod{1}$ and $C(i + j + 1, j + 1)\beta \pmod{1}$ are in the interval $(1/8, 3/8)$, while $[C(i + j + 1, j) + C(i + j, j + 1)]\beta \pmod{1}$ is in the interval $(3/16, 9/16)$. Continue, making it impossible that $\lambda^{r_n} \rightarrow 1$. \square

5. CONDITION FOR TOPOLOGICAL CONJUGACY TO AN ODOMETER

According to the theorem of Downarowicz and Maass [6] mentioned above, every proper minimal adic system with finite topological rank is either expansive or topologically conjugate to an odometer. For systems with a bounded number of vertices per level some sufficient conditions for topological conjugacy to an odometer are known (see the remarks in the introduction). In this section we give a necessary and sufficient condition for a fairly general adic system to be topologically conjugate to an odometer. The condition is then used to determine when, with probability 1, an adic system with a random ordering, or an adic system constructed according to a random process, is topologically conjugate to an odometer.

In this section we deal with an arbitrary Bratteli diagram $B = (\mathcal{V}, \mathcal{E})$. As usual the vertices are arranged into levels, so that $\mathcal{V} = \cup_{i \geq 0} \mathcal{V}_i$, $\mathcal{E} = \cup_{i \geq 1} \mathcal{E}_i$, and there are directed

edges from each level to the next: there are a range map r and a source map s from \mathcal{E} to \mathcal{V} such that $r(\mathcal{E}_i) = \mathcal{V}_i$ for all $i \geq 1$, $s(\mathcal{E}_i) = \mathcal{V}_{i-1}$ for all $i \geq 1$, $s^{-1}v \neq \emptyset$ for all $v \in \mathcal{V}$, and $r^{-1}v' \neq \emptyset$ for all $v' \in \mathcal{V} \setminus \mathcal{V}_0$. We assume that \mathcal{V}_0 consists of a single vertex, v_0 , and $|\mathcal{V}_n|, |\mathcal{E}_n| < \infty$ for all n .

The *path space* X is the space of infinite paths $x = x_1x_2\dots$ such that $x_i \in \mathcal{E}_i$ and $r(x_i) = s(x_{i+1})$ for all i . To avoid degenerate cases, we assume that X with the usual metric (in which two infinite paths are close if they agree on long initial segments) is homeomorphic to the Cantor set. An *ordering* ξ of the diagram is an assignment of a linear order to the set $r^{-1}v$ of edges entering each vertex $v \in \mathcal{V}$. $X_{\xi, \max}$ consists of the set of paths all of whose edges are maximal in the ordering ξ , and $X_{\xi, \min}$ of all the paths all of whose edges are minimal. As with the Pascal diagram, the ordering ξ determines the adic transformation $T_\xi : X \setminus X_{\xi, \max} \rightarrow X \setminus X_{\xi, \min}$, which is a homeomorphism.

See [15], [12], or [7] for the definitions of *telescoping* and *isomorphism* for diagrams and ordered diagrams. *Equivalence* of diagrams or ordered diagrams is the equivalence relation generated by isomorphism and telescoping. Two diagrams B_1 and B_2 are diagram equivalent if and only if there is a diagram B such that telescoping B to odd levels yields a telescoping of one of B_1, B_2 and telescoping B to even levels yields a telescoping of the other one; similarly for ordered diagrams. If two ordered diagrams are equivalent, then the topological dynamical systems that they define by means of their adic transformations are topologically conjugate. Recall that a Bratteli diagram is called *simple* if there is a telescoping for which every pair of vertices $v_n \in \mathcal{V}_n$ and $v_{n+1} \in \mathcal{V}_{n+1}$ there is an edge between v_n and v_{n+1} . That the converse holds also, for simple adic systems that are properly ordered (with unique maximal and minimal paths), follows from Theorem 4.4 of [15]. (For suppose that simple and properly ordered diagrams D_1 and D_2 determine adic systems X_1 and X_2 , respectively, that are topologically conjugate under an adic-commuting homeomorphism $h : X_1 \rightarrow X_2$. Each X_i has a natural sequence of Kakutani-Rokhlin partitions \mathcal{P}_i corresponding to its diagram structure. Then $h^{-1}\mathcal{P}_2$ is a sequence of Kakutani-Rokhlin partitions for X_1 that is admissible as in Theorem 4.2 of [15]. This sequence determines D_2 according to the process in [15], so by Theorem 4.4 of [15] D_1 and D_2 are diagram equivalent.) (Note that in the proof of Theorem 4.4 of [15] the tops Z_n and \tilde{Z}_n of the tower diagrams have the property that given n we can find m such that \tilde{Z}_m is contained in Z_n , and vice versa.)

An *odometer* is any topological dynamical system that is topologically conjugate to the system defined by the adic transformation on the path space of an ordered diagram which has exactly one vertex at every level.

For each n and each $w \in \mathcal{V}_n$, there is an associated coding in terms of vertices in \mathcal{V}_{n-1} . (This is the ‘‘morphism read on \mathcal{V}_n ’’ in [7, p. 342].) More precisely, if $r^{-1}w =$




FIGURE 10. A uniformly ordered level, where $\omega = v_2v_1v_3$.

$\{e_1, e_2, \dots, e_{|r^{-1}w|}\}$, where the subscripts correspond directly to the ordering given by ξ , then the coding of w by vertices in \mathcal{V}_{n-1} is given by $c(w) = s(e_1)s(e_2) \dots s(e_{|r^{-1}w|})$. We call \mathcal{E}_n and the level n *uniformly ordered* if there is a string $v = v_{i_1}v_{i_2} \dots v_{i_k}$, with each v_{i_m} in V_{n-1} , such that for every w in \mathcal{V}_n , $c(w) = v^{k_w}$ for some $k_w \in \mathbb{N}$. See Figure 10. Note that if any level has only one vertex, then it is automatically uniformly ordered.

For a Bratteli-Vershik system $B = (\mathcal{V}, \mathcal{E})$, define an *ordered shape at level n* to consist of \mathcal{V}_{n-1} , \mathcal{V}_n , \mathcal{E}_n , and a partial order on \mathcal{E}_n which is a total order on each $r^{-1}w, w \in \mathcal{V}_n$. We say that a shape is *uniformly ordered* if \mathcal{E}_n is uniformly ordered. Note that each row of the incidence matrix that corresponds to a uniformly ordered shape is an integer multiple of a single vector.

Lemma 5.1. *If a diagram $B = (\mathcal{V}, \mathcal{E})$ with an ordering ξ has infinitely many levels for which \mathcal{E}_n is uniformly ordered, then (1) there is a telescoping $B' = (\mathcal{V}', \mathcal{E}')$ of $B = (\mathcal{V}, \mathcal{E})$ for which every level is uniformly ordered, and (2) $B = (\mathcal{V}, \mathcal{E})$ is simple and properly ordered.*

Proof. First note that if an ordered Bratteli diagram is telescoped from level j_1 to level j_2 , the new coding of the vertices at level j_2 in terms of the vertices at level j_1 results from decomposing the original $j_2 - 1$ coding of vertices in \mathcal{V}_{j_2} by vertices in \mathcal{V}_{j_2-1} in an inductive manner into a level j_1 coding. Then note that if $j_1 < n \leq j_2$ and \mathcal{E}_n is uniformly ordered, then the new set of edges arising from telescoping from level j_1 to level j_2 will be uniformly ordered. Indeed, since all vertices $w \in \mathcal{V}_n$ have codings $c(w) = v^{k_w}$ in terms of the vertices at level $n - 1$, decomposing the level $j_2 - 1$ codings of vertices in \mathcal{V}_{j_2} into level $n - 1$ codings will lead to repeated concatenations of v . Furthermore, the ordering ξ will lead to a unique decomposition of v in terms of vertices at level j_1 , and therefore the new set of edges will be uniformly ordered. See Figure 11.

In order for a level to be uniformly ordered, every vertex in \mathcal{V}_n must connect to every vertex in \mathcal{V}_{n+1} . Hence any diagram with infinitely many uniformly ordered levels is simple.

For every n there is a unique vertex v_{n-1} in \mathcal{V}_{n-1} such that every edge in \mathcal{E}_n which is maximal has v_{n-1} as its source. Therefore any path in X_{\max} must go through these vertices, and there is a unique path which travels along maximal edges at every level. \square




FIGURE 11. Following the code shows that it is possible to telescope so that every level is uniformly ordered.

Remark 5.2. It is possible for two non-uniformly ordered levels to telescope to one that is uniformly ordered. Indeed, suppose that $|\mathcal{V}_n| \geq 3$ and the level $n - 1$ coding of vertices at level n is such that $c(w_2) = c(w_3) \neq c(w_1)$ for w_1, w_2, w_3 in \mathcal{V}_n . Then suppose that the level n coding of vertices in \mathcal{V}_{n+1} is such that the only difference in the codings of different vertices is that w_2 and w_3 are interchanged in two codings. Then neither \mathcal{E}_n nor \mathcal{E}_{n+1} is uniformly ordered. However, once one telescopes between levels $n - 1$ and $n + 1$, the new set of edges will be uniformly ordered. See Figure 12




FIGURE 12. All orderings are left to right. Neither level is uniformly ordered, but the telescoping is.

Theorem 5.3. *If an ordered Bratteli diagram $B = (\mathcal{V}, \mathcal{E}, \xi)$ has a telescoping for which there are infinitely many uniformly ordered levels then the associated adic system is topologically conjugate to an odometer. Conversely, any simple properly ordered Bratteli diagram $B = (\mathcal{V}, \mathcal{E}, \xi)$ that is topologically conjugate to an odometer has a telescoping for which there are infinitely many uniformly ordered levels.*

Proof. Suppose that we have a diagram $B = (\mathcal{V}, \mathcal{E}, \xi)$ for which there is a telescoping with infinitely many uniformly ordered levels. By Lemma 5.1 we may assume that the diagram is uniformly ordered at every level $n > 1$; also note that the diagram is simple and properly ordered. One can see then that all the k -codings (coding T_ξ -orbits according to the initial path segments of length k) are periodic, so that the system is not expansive and hence, by [6], is topologically conjugate to an odometer—in case the system has finite topological rank. We give here an explicit proof in the general case by constructing an intermediate ordered Bratteli diagram, B' , with the property that telescoping to odd levels yields the given diagram, B , and telescoping to even levels yields a system with a single vertex at every level. Thus B is equivalent to the diagram of an odometer, and hence its associated topological dynamical system is topologically conjugate to an odometer.

To construct B' , for each $n \geq 1$ let $|\mathcal{V}'_{2n-1}| = |\mathcal{V}_n|$ and $|\mathcal{V}'_{2n}| = 1$. Let $\mathcal{E}'_1 = \mathcal{E}_1$. For every $n \geq 1$, denote the single vertex in \mathcal{V}'_{2n} by w'_{2n} and set up a one-to-one correspondence between \mathcal{V}'_{2n-1} and \mathcal{V}_n . Fix $n \geq 1$. Since B is uniformly ordered at level $n + 1$, there is a string $v = v_1 \dots v_{|v|}$ with entries from \mathcal{V}_n such that for each w in \mathcal{V}_{n+1} the coding of w by vertices in \mathcal{V}_n is $c(w) = v^{k_w}$. To define \mathcal{E}'_{2n} let $|r^{-1}w'_{2n}| = |v|$, label the edges e'_i in $r^{-1}w'_{2n}$ so that for each i $s(e'_i)$ is the vertex u' in \mathcal{V}'_{2n-1} which corresponds to v_i , and order these edges according to their labels. To define \mathcal{E}'_{2n+1} , note that each $u' \in \mathcal{V}'_{2n+1}$ corresponds to a $w \in \mathcal{V}_{n+1}$; we let the number of edges from w'_{2n} to u' be k_w for the w corresponding to u' and order these edges arbitrarily. Then telescoping B' to odd levels yields B , and telescoping to even levels yields a system with just one vertex at every level, an odometer. See Figure 13.

Conversely, suppose that $B = (\mathcal{V}, \mathcal{E}, \xi)$ is a simple properly ordered diagram whose associated adic system is topologically conjugate to an odometer. As mentioned above, then the diagram is equivalent to an odometer: there is an intermediate diagram $B' = (\mathcal{V}', \mathcal{E}', \xi')$, which when telescoped to odd levels yields a telescoping B'' of B and when telescoped to even levels yields a system with just one vertex at each level, or vice versa. For each n , $|\mathcal{V}'_{2n}| = 1$ and hence level $2n$ of B' is automatically uniformly ordered. Then as in the proof of Lemma 5.1, the telescoping of B' from level $2n - 1$ to level $2n + 1$ will yield a set of edges whose ranges form a uniformly ordered level. Therefore in B'' every level \mathcal{E}_n must be uniformly ordered. \square




FIGURE 13. B' telescopes to B on odd levels, and to the odometer on even levels. Give B any uniform ordering and the diagrams are order equivalent.

We are seeking conditions that will allow one to determine when BV systems are expansive or not, conjugate to odometers or not, and of finite topological rank or not. As mentioned above, Downarowicz and Maass [6] (see also [17]) showed that every properly ordered minimal BV system of finite topological rank is either expansive or else conjugate to an odometer. Gjerde and Johansen [13] give an example of a simple properly ordered BV system that is not expansive, yet not conjugate to an odometer, thus necessarily of infinite topological rank. We note in the following Proposition that the key properties of this example are enjoyed by a wide class of systems, which we call “standard nonexpansive”, and which therefore are nonexpansive and, by essentially the same argument as in [13], not of finite topological rank, in particular not conjugate to odometers.

Recall that given $N \in \mathbb{N}$, every path in an ordered BV system $B = (\mathcal{V}, \mathcal{E}, \xi)$ has a N -coding of its orbit according to the partition \mathcal{P}_N into cylinder sets to level N . For each vertex $v \in \mathcal{V}$, the edges with range v are assigned labels indicating their positions in the ordering determined by ξ . So any path γ has an infinite *sequence of edge labels* associated with it. Specifically, the i th entry in the sequence is the label assigned to the i th edge of γ . We say that the BV system is *standard nonexpansive* if for every natural number N there exist distinct paths γ_N and γ'_N that are assigned the same sequence of edge labels, agree from the root to some level $K_N > N$, and at any level the vertices met by the two paths have the same basic block in the N -coding. In this case, γ_N and γ'_N have the same N -coding. Standard nonexpansive implies nonexpansive.

Proposition 5.4. *Every standard nonexpansive simple properly ordered BV system has infinite rank.*

Proof. By [15], for any BV system B that is conjugate to the odometer there is a BV system B' that telescopes on odd levels to a telescoping of B and on even levels to a telescoping of the odometer. There is then a telescoping of B in which every vertex is uniformly ordered.

This, however, cannot occur when there are paths with the same sequence of edge labels. So if B is standard nonexpansive, then it not conjugate to the odometer. It then follows, by [6], that B does not have finite topological rank. \square

We consider now what might happen if a diagram is ordered randomly, or if a diagram is somehow chosen at random.

Suppose that we are given a simple diagram $B = (\mathcal{V}, \mathcal{E})$ as above. Denote the set of orderings of the edges entering a vertex $v \in \mathcal{V}$ by \mathcal{O}_v . The set of all orderings of B is $\mathcal{O} = \prod_{v \in \mathcal{V}} \mathcal{O}_v$. We consider probability measures μ on \mathcal{O} , for example the uniform and independent measure μ_0 which is the product over all v of the uniform measures on the discrete spaces \mathcal{O}_v . For each $n = 1, 2, \dots$ let U_n denote the set of orderings of B which are uniformly ordered at level n , and let U denote the set of orderings which have infinitely many uniformly ordered levels. By the preceding theorem, if $\mu(U) = 1$ then an ordering selected randomly according to μ will with probability 1 produce an adic system topologically conjugate to an odometer.

Example 5.5. Suppose that B is a diagram such that for every $n \geq 1$ there is an r_n such that the number of edges between every pair of vertices $v_{n-1} \in \mathcal{V}_{n-1}$ and $v_n \in \mathcal{V}_n$ is r_n . (Then B is simple.) Write $|\mathcal{V}_n| = V_n$. Then there are $r_n!$ orderings of the edges into each vertex at level n and thus $(r_n!)^{V_n}$ ways to order the unordered shape at level n . There are $r_n!$ uniform orderings at level n , since we must have the same ordering at each vertex on a fixed level. By the Borel-Cantelli Lemma, if

$$(6) \quad \sum_{n=1}^{\infty} \frac{r_n!}{(r_n!)^{V_n}} = \infty,$$

then $\mu_0(U) = 1$, and so with probability 1 the system is topologically conjugate to an odometer. For example, if the r_n and V_n are bounded, and B is as described, then selecting orderings uniformly and independently yields an odometer with probability 1. This is consistent with Corollary 5.3 of [4], according to which a diagram with bounded number of vertices and edges per level will have a “perfect” ordering (an ordering which admits a Vershik map as a homeomorphism) with probability 1.

Example 5.6. We can also consider assembling a random ordered diagram from a sequence of shapes, that fit together, generated by a stochastic process. Generalizing the definition above, define an *ordered shape* $S = (C, D, E, \xi)$ to be a bipartite directed graph together with a partial order on its set of edges such that (1) the vertex set is the disjoint union of two finite sets C and D ; (2) for each $e \in E$, $s(e) \in C$ and $r(e) \in D$; (3) for each $b \in D$, ξ is a total order on $r^{-1}b$; (4) for $b_1, b_2 \in D$ with $b_1 \neq b_2$, the edges in $r^{-1}b_1$ and $r^{-1}b_2$ are not comparable.

Let \mathcal{S} be the set of sequences of ordered shapes $S_n = (C(S_n), D(S_n), E(S_n), \xi(S_n))$, $n = 1, 2, \dots$, such that $C(S_{n+1}) = D(S_n)$ for all n , so that the shapes can “fit together”. If μ is a probability measure on \mathcal{S} that assigns full measure to the set of sequences which have infinitely many uniformly ordered shapes, then constructing an adic system at random with respect to μ produces an odometer with probability 1. For example, if μ is an ergodic shift-invariant probability measure on \mathcal{S} such that $\mu\{S = (S_n) \in \mathcal{S} : S_1 \text{ is uniformly ordered}\} > 0$, then by the Poincaré Recurrence Theorem S will have infinitely many uniformly ordered shapes and hence will describe an odometer.

6. THE UNION OVER ALL ORDERINGS

For a subshift (Ω, σ) , denote by $\mathcal{L}(\Omega)$ the *language* of Ω , that is, the set of finite blocks seen in sequences in Ω . Define for each ordering ξ of the Pascal graph P the subshift $\Sigma_\xi \subset \{a, b\}^{\mathbb{Z}}$ to consist of all doubly infinite sequences, each of whose subblocks can be found in the 1-coding of some $\gamma \in X$ (the set of infinite paths on the Pascal graph) for the ordering ξ , and define the “big subshift” (Σ, σ) by $\mathcal{L}(\Sigma) = \cup_\xi \mathcal{L}(\Sigma_\xi)$. Equivalently, we can define the big subshift Σ to be the closure of the union over all orderings ξ of X_ξ :

$$(7) \quad \Sigma = \overline{\bigcup_{\xi} X_{\xi}}.$$

What is the asymptotic complexity of Σ ? What are other dynamical properties of (Σ, σ) ? Is this system topologically transitive (i.e., does it have a dense orbit)? Are there periodic points?

Recall the basic blocks $B_\xi(x, y)$ defined above. Basic blocks are determined by the orderings on the edges above them. See Figure 14. However, a block may be generated by




FIGURE 14. Each choice of ordering of the incoming edges to the vertices in the shaded region will determine a $B_\xi(x, y)$.




FIGURE 15. Dashed lines represent minimal edges. Both orderings ξ_1 and ξ_2 give rise to $B(4, 2) = a^2ba^2ba^2b^2a^2ba^2$.

different orderings. For example, there exist two distinct orderings, ξ_1 and ξ_2 such that $B_{\xi_1}(2, 1) = B_{\xi_2}(2, 1) = aba$. This can continue throughout the diagrams. See Figure 15.

Assuming a fixed ordering ξ , the basic blocks $B(x, 1)$, for $x \geq 1$, will be used frequently in the following and we will give them the special name C_x . In a similar fashion, for $y \geq 1$ define the block $D_y = B(1, y)$.

In the following we will consider the set of diagrams for which

$$\xi(x, 1) = \xi(1, y) = 0.$$

In other words, the ordering on the edges coming into vertices $(x, 1)$ and $(1, y)$ is left to right for all values of $x \geq 1$ and $y \geq 1$. This means that $C_i = a^i b$ and $D_j = ab^j$ for $i, j \geq 1$. In the following we will show that for such orderings basic blocks have unique factorizations.

Lemma 6.1. *Given (x, y) with $x \geq 1$ and $y \geq 1$, there is a one-to-one correspondence between the set of possible basic blocks*

$$\{B_\xi(x, y) : \xi(i, 1) = \xi(1, j) = 0 \text{ for all } i, j \geq 1\}$$

and the set of orderings of edges into each vertex (u, v) with $u \leq x$ and $v \leq y$: $B_\xi(x, y) = B_\eta(x, y)$ if and only if $\xi(u, v) = \eta(u, v)$ for all (u, v) with $u \leq x$ and $v \leq y$.

Proof. The reverse direction is clear from the definition of $B_\xi(x, y)$. We will therefore consider a $B_\xi(x, y)$ and show that it gives rise to a unique ξ , given the appropriate restrictions.

Assume $x, y > 1$ and we are given a basic block $B_\xi(x, y)$. Starting from the left in $B_\xi(x, y)$, find the first occurrence of a block of the form, for some $i_0, j_0 \geq 1$, $a^{i_0} b^{j_0} a$, or $a^{i_0} b^{j_0}$ if no $a^{i_0} b^{j_0} a$ is to be found. Because of the restriction on ξ , we know that it is either the case that $i_0 = 1$ and $j_0 \neq 1$ or $i_0 \neq 1$ and $j_0 = 1$. In the first case, $a^{i_0} b^{j_0}$ corresponds to the block D_{j_0} , and in the second to C_{i_0} . Continue left to right in this fashion to decompose $B_\xi(x, y)$ completely into a sequence of C_i 's and D_j 's. Note that there is a unique path

between $(x, 1)$ and (x, y) , and therefore C_x appears only once in $B_\xi(x, y)$. Furthermore, if $i > x$ then there is no path from $(i, 1)$ to (x, y) , and therefore C_i cannot be a subblock of $B(x, y)$. A similar argument gives the result for the D_y . So C_x and D_y each appear exactly once, and if C_i (or D_j) appears in $B_\xi(x, y)$, then $i \leq x$ (or $j \leq y$).

Now, $B_\xi(x, y)$ decomposes into the concatenation of $B_\xi(x - 1, y)$ and $B_\xi(x, y - 1)$. Note that the unique C_x in $B_\xi(x, y)$ is a subblock of the $B_\xi(x, y - 1)$ portion of $B_\xi(x, y)$. Likewise, D_y is a subblock of the $B_\xi(x - 1, y)$ portion of $B_\xi(x, y)$. Therefore, if C_x appears before D_y , then $B_\xi(x, y) = B_\xi(x, y - 1)B_\xi(x - 1, y)$, and vice versa. This determines the order that ξ gives to the edges with range (x, y) . We also know the lengths of each of $B(x, y - 1)$ and $B(x - 1, y)$. Once we know which one comes first, we know exactly where the juxtaposition between the two occurs. We repeat this process on each word above $B(x, y)$ to conclude that no ordering besides ξ could have produced the block $B_\xi(x, y)$. \square

We work through an example.

Example 6.2. Consider the following block:

$$ab^3ab^2a^2ba^3bab^2a^2ba^3bab^2a^2ba^4b$$

Because we know the structure of the C_i and D_j , we see that the first set of a 's and b 's is ab^3 . This must be D_3 . Then the next set of a 's and b 's is ab^2 . This must be D_2 . We continue in this manner to see that the above block decomposes into $D_3D_2C_2C_3D_2C_2C_3D_2C_2C_4$. By the above lemma, this must be block $B(4, 3)$. Because the D_3 appears before the C_4 , $B(4, 3)$ must decompose into $B(3, 3)B(4, 2)$, which implies there was a right to left ordering on the edges with range $(4, 3)$. We need only to know that $|B(3, 3)| = 20$ to see that $B(3, 3) = D_3D_2C_2C_3D_2C_2$ and $B(4, 2) = C_3D_2C_2C_4$. We continue in this manner until we have placed an order on the incoming edges of all 6 unordered vertices that are above $(4, 3)$. See Figure 16.




FIGURE 16. The ordering of edges determined by $B(4, 3) = ab^3ab^2a^2ba^3bab^2a^2ba^3bab^2a^2ba^4b$

Proposition 6.3. *Given left to right ordering on edges terminating at vertices $(u, 1)$ and $(1, v)$ for all $u, v \geq 1$, then for all $x, y \geq 1$ there are $2^{(x-1)(y-1)}$ distinct blocks $B_\xi(x, y)$ as we vary the orderings ξ .*

Proof. By the Lemma, each ordering ξ of the edges entering vertices (u, v) above (x, y) ($1 \leq u \leq x, 1 \leq v \leq y$) gives rise to a distinct $B_\xi(x, y)$. There are $(x-1)(y-1)$ such vertices (u, v) above (x, y) and two choices of ordering for each vertex, yielding $2^{(x-1)(y-1)}$ distinct orderings and hence that many blocks $B_\xi(x, y)$. \square

Proposition 6.4. *For the big subshift, the complexity function $p(n)$ is superpolynomial: for each $k = 2, 3, \dots$ there are at least 2^{k-1} blocks of length $(k+1)(k+2)/2$ in the big subshift.*

Proof. Let $k = 2, 3, \dots$ and $n = (k+1)(k+2)/2$. Note that the binomial coefficient $C(k+2, 2) = n$. Then for any ordering ξ , $B_\xi(k, 2)$ has length n since the length of $B(k, 2)$ is the binomial coefficient $C(k+2, 2)$.

Then the coarse lower bound above is given by just looking at the 2^{k-1} unique ways to form $B_\xi(k, 2)$ and applying Proposition 6.3. \square

We will show next that the big subshift Σ is not topologically transitive and that the “small subshift” whose language is $\cap_\xi \mathcal{L}(\Sigma_\xi)$ consists of only the two fixed points and the orbits of the points $a^\infty.ba^\infty$ and $b^\infty.ab^\infty$.

Lemma 6.5. (1) *There are orderings ξ and ξ' of the Pascal graph such that $B_\xi(3, 3) = a(ab)^9b$ and $B_{\xi'}(3, 3) = b(ba)^9a$. (2) *For every ordering ξ and every $x, y \geq 0$, the basic block $B_\xi(x, y)$ does not contain both $(ab)^9$ and $(ba)^9$.**

Proof. To prove (1), we proceed to define the orderings ξ and ξ' called for in the Lemma. Define $\xi(1, 1) = \xi'(1, 1) = 0$, so that $B_\xi(1, 1) = B_{\xi'}(1, 1) = ab$. Further, choose ξ and ξ' so that $B_\xi(2, 1) = B_{\xi'}(2, 1) = aba$ and $B_\xi(1, 2) = B_{\xi'}(1, 2) = bab$. Continuing, define ξ and ξ' so that $B_\xi(3, 1) = a^2ba$, $B_\xi(2, 2) = (ba)^3$, and $B_\xi(1, 3) = bab^2$, while $B_{\xi'}(3, 1) = aba^2$, $B_{\xi'}(2, 2) = (ab)^3$, and $B_{\xi'}(1, 3) = b^2ab$. Then define $\xi(3, 2) = \xi(2, 3) = \xi(3, 3) = 0$ (so that the ordering by ξ is “left-right” at these vertices) and $\xi'(3, 2) = \xi'(2, 3) = \xi'(3, 3) = 1$ (so that the ordering by ξ' is “right-left” at these vertices). We then have $B_\xi(3, 3) = a(ab)^9b$ and $B_{\xi'}(3, 3) = b(ba)^9a$. See Figure 17.

To prove (2), we want to focus on strings of alternating a 's and b 's that appear in the language of Σ_ξ for some ordering ξ of the Pascal graph and hence in a basic block $B_\xi(x, y)$ for some $x, y \geq 0$. So we provide a systematic way to ignore subblocks of the $B_\xi(x, y)$ that cannot be used, when $B_\xi(x, y)$ is concatenated with $B_\xi(x-1, y+1)$ or $B_\xi(x+1, y-1)$, to




FIGURE 17. Dashed lines represent minimal edges. $B_\xi(3, 3) = a(ab)^9b$ and $B_{\xi'} = b(ba)^9a$.

create a longer string $(ab)^j$ than has been created previously. Specifically, for each $x, y \geq 1$ we let $Q_\xi^L(x, y)$ (respectively $Q_\xi^R(x, y)$) be the longest string in which a and b alternate that is a prefix of $B_\xi(x, y)$ (respectively suffix of $B_\xi(x, y)$). For example, if $B_\xi(x, y) = ababa^2b$, then $Q_\xi^L(x, y) = ababa$ and $Q_\xi^R(x, y) = ab$. We add the new symbol $*$ to our alphabet and define the “condensed form” of $B_\xi(x, y)$ to be $Q_\xi(x, y) = Q_\xi^L(x, y) * Q_\xi^R(x, y)$. (If the block $B_\xi(x, y)$ is made up entirely of alternating a ’s and b ’s there is no $*$ in $Q_\xi(x, y)$.)

We want to show that if ξ is an ordering such that $(ab)^9 \in \mathcal{L}(\Sigma_\xi)$, then $(ba)^9 \notin \mathcal{L}(\Sigma_\xi)$, and vice versa; that is, no $B_\xi(x, y)$ can ever contain both $(ab)^9$ and $(ba)^9$. We begin with two observations. The first is that for $x \geq 3$, $Q_\xi^L(x, 1)$ cannot have length greater than 3. (Specifically, $Q_\xi^L(x, 1)$ can be aba but cannot be any longer.) The same is true for $Q_\xi^R(x, 1)$, and similarly for $Q_\xi^L(1, y)$ and $Q_\xi^R(1, y)$ for $y \geq 3$. The second observation is that if a basic block B_1 has condensed form $C_1^L * C_1^R$ and a second basic block B_2 has condensed form $C_2^L * C_2^R$, then the concatenation B_1B_2 has condensed form $C_1^L * C_2^R$. Therefore, in order to determine the highest power of $(ab)^j$ that can be formed by a concatenation of any two basic blocks one needs to look only far enough down to a level n for which the condensed forms of all the basic blocks $B_\xi(x, y)$ with $x + y = n$ and $x, y \geq 1$ contain the symbol $*$. We can then determine the highest power of $(ab)^j$ in $\mathcal{L}(\Sigma_\xi)$ by considering all possible concatenations $Q_\xi^R(i, j)Q_\xi^L(k, l)$ with $i + j = k + l = n$ and $i, j, k, l \geq 1$.

Without loss of generality we assume that $B_\xi(1, 1) = ab$. Then depending on the choice among the four partial orders on the edges with range in \mathcal{V}_3 there are the following possibilities for the basic blocks at level 3: $B_\xi(3, 0) = a$, $B_\xi(2, 1) = a^2b$ or aba , $B_\xi(1, 2) = ab^2$ or bab and $B_\xi(0, 3) = b$. It is straightforward to check that for three of the four possible orderings all the condensed blocks $Q_\xi(x, y)$ with $x + y = 4$ and $x, y \geq 1$ necessarily contain the symbol $*$. Since every basic block at level 4 has length no more than 6, for these three orderings it is impossible that $(ab)^9$ or $(ba)^9$ is contained within any of these blocks or, since they all contain a $*$, any of their concatenations. Hence the only way that $(ab)^9$ or $(ba)^9$ could be formed is in the case of the fourth ordering at level 3, when $B_\xi(2, 1) = aba$ and $B_\xi(1, 2) = bab$. In this case the possibilities are $Q_\xi(3, 1) = a*aba$ or $aba*a$, $Q_\xi(2, 2) = (ab)^3$

or $(ba)^3$, and $Q_\xi(1, 3) = bab * b$ or $b * bab$. For each of the 8 possible orderings at the vertices (x, y) with $x + y = 5$ and $x, y \geq 1$ the condensed blocks $Q_\xi(x, y)$ all necessarily contain a $*$. Therefore 5 is the last level at which we need to look in order to determine which orderings give rise to either $(ab)^9$ or $(ba)^9$.

The only two blocks on level 5 that could concatenate to give a word of length 18 are $B_\xi(3, 2)$ and $B_\xi(2, 3)$, both of length 10. Further, since each of $Q_\xi(3, 2)$ and $Q_\xi(2, 3)$ contains a $*$, the maximum length of $Q_\xi^L(3, 2), Q_\xi^R(3, 2), Q_\xi^L(2, 3), Q_\xi^R(2, 3)$ is 9. In order to form either $(ab)^9$ or $(ba)^9$, we would need both $Q_\xi^L(3, 2)$ and $Q_\xi^R(2, 3)$ to be of length 9, or both $Q_\xi^R(3, 2)$ and $Q_\xi^L(2, 3)$ to be of length 9. In the situation to which we have reduced, there is exactly one ordering down to level 5 for which both $Q_\xi^L(3, 2)$ and $Q_\xi^R(2, 3)$ have length 9; for this ordering $Q_\xi(3, 2) = (ab)^4 a * a$ and $Q_\xi(2, 3) = b * b(ab)^4$. In this case it is possible to concatenate $B_\xi(3, 2)$ and $B_\xi(2, 3)$ to form the word $(ba)^9$. However, because these two blocks contain the longest possible strings of alternating a 's and b 's down to level 5, there is no way, given this ordering, ever to create the word $(ab)^9$. Similarly, there is exactly one case for which $Q_\xi^R(3, 2)$ and $Q_\xi^L(2, 3)$ are of length 9, namely when $Q_\xi^R(3, 2) = a * a(ba)^4$ and $Q_\xi^L(2, 3) = (ba)^4 b * b$. The string $(ab)^9$ can be made with the ordering corresponding to this case, but $(ba)^9$ cannot. Therefore, any ordering which gives rise to the block $(ab)^9$ cannot also give rise to $(ba)^9$, and vice versa. \square

Theorem 6.6. *There is no dense orbit in Σ .*

Proof. By the Lemma, both of the blocks $(ab)^9$ and $(ba)^9$ are in $\mathcal{L}(\Sigma) = \cup_\xi \mathcal{L}(\Sigma_\xi)$. But no sequence in Σ can contain both of these blocks, since any block containing both of them would have to be a subblock of $B_\xi(x, y)$ for some ξ, x , and y . \square

Proposition 6.7. *The only periodic points in the big subshift (Σ, σ) are the two fixed points a^∞ and b^∞ .*

Proof. Suppose that ω is a periodic sequence. If ω is not one of the two fixed points, then the runs of a and b in ω are of bounded length: there is $r < \infty$ such that if a^j or b^j appear in ω , then $j < r$. For each ordering ξ the two basic blocks $B_\xi(2r, 1)$ and $B_\xi(1, 2r)$ each contain strings of length r of a or b , respectively. If $n > 4r$, then every basic block $B_\xi(x, y)$ with $x + y = n$ and $x, y \geq 1$, which is a concatenation of basic blocks at level $2r + 1$, contains one of the basic blocks $B_\xi(1, 2r)$ or $B_\xi(2r, 1)$ from level $2r + 1$. Denote by M the maximum of the lengths of basic blocks at level $4r$. Every block in the language $\mathcal{L}(\Sigma)$ of length greater than $3M$ must contain either a basic block of length greater than r from level $4r$, and hence a^j or b^j for some $j \geq r$, or one of the strings a^M or b^M . The later cannot occur since $M > r$. Thus no block in ω of length greater than $3M$ can be in $\mathcal{L}(\Sigma)$, and ω cannot be in Σ unless it is one of the two fixed points. \square

Lemma 6.8. *The small subshift $\cap_{\xi} \Sigma_{\xi}$ contains the points $a^{\infty}, b^{\infty}, a^{\infty}.ba^{\infty}$, and $b^{\infty}.ab^{\infty}$.*

Proof. If $\xi(x, 1)$ is not eventually constant, then for every $r > 0$ there is an x such that $B_{\xi}(x, 1) = a^i b a^j$ and $i, j > r$. Then $a^{\infty}.ba^{\infty}$ and a^{∞} are in Σ_{ξ} .

Suppose then that $\xi(x, 1)$ is eventually constant, say $\xi(x, 1) = 0$ for all $x > K$. Then $B_{\xi}(x, 1) = a^{x-m} b a^m$ for some $m \leq x$ and all $x > K$. The block $B_{\xi}(x, 2)$ is a concatenation of the blocks $B_{\xi}(x-3, 2)$, $B_{\xi}(x-2, 1)$, $B_{\xi}(x-1, 1)$ and $B_{\xi}(x, 1)$ in some order, each with multiplicity one. By the pigeonhole principle it is necessary that either $B_{\xi}(x-2, 1)$ is concatenated with $B_{\xi}(x-1, 1)$, or $B_{\xi}(x-2, 1)$ is concatenated with $B_{\xi}(x, 1)$, or $B_{\xi}(x-1, 1)$ is concatenated with $B_{\xi}(x, 1)$. It follows that for $x > K+3$ the block $a^{x-m-2} b a^{x-2}$ necessarily appears in $B_{\xi}(x, 2)$. Letting x tend to infinity shows that $a^{\infty}.ba^{\infty}$ and a^{∞} are in Σ_{ξ} .

That $b^{\infty}.ab^{\infty}, b^{\infty} \in \Sigma_{\xi}$ follows by symmetry. \square

Theorem 6.9. *The small subshift $\cap_{\xi} \Sigma_{\xi}$ consists of only the two fixed points and the orbits of the points $a^{\infty}.ba^{\infty}$ and $b^{\infty}.ab^{\infty}$.*

Proof. We fix the orderings ξ and ξ' used in Lemma 6.5:

$$(8) \quad \begin{aligned} \xi(1, 1) = \xi'(1, 1) = 0, \xi(1, 2) = \xi'(1, 2) = \xi(2, 1) = \xi'(2, 1) = 1, \\ \xi(2, 2) = 1, \xi'(2, 2) = 0; \xi(1, 3) = \xi(3, 1) = 0, \xi'(1, 3) = \xi'(3, 1) = 1 \\ \text{for all } (x, y) \text{ with } x + y \geq 5, \xi(x, y) = 0; \\ \text{for all } (x, y) \text{ with } x + y \geq 5, \xi'(x, y) = 1. \end{aligned}$$

For notational ease, let $B_{\xi}(j, 2)$ be denoted by F_j , let $B_{\xi}(2, k)$ be denoted by G_k , and let $B_{\xi}(3, 3) = a(ab)^9 b = H$. Likewise, let $B_{\xi'}(j, 2)$ be denoted by F'_j , let $B_{\xi'}(2, k)$ be denoted by G'_k , and let $B_{\xi'}(3, 3) = b(ba)^9 a = H'$. For all $j, k \geq 4$,

$$(9) \quad \begin{aligned} F_j = B_{\xi}(j, 2) &= a^{j-1} b a^{j-1} b a^{j-2} b a^{j-3} b \dots a^4 b a^3 (b a)^4 \\ G_k = B_{\xi}(2, k) &= (b a)^4 b^3 a b^4 a b^5 a \dots b^{k-1} a b^{k-1} \end{aligned}$$

and

$$(10) \quad \begin{aligned} F'_j = B_{\xi'}(j, 2) &= (a b)^4 a^3 b a^4 b a^5 b \dots b a^{j-1} b a^{j-1} \\ G'_k = B_{\xi'}(2, k) &= b^{k-1} a b^{k-1} a b^{k-2} a b^{k-3} \dots b^5 a b^4 a b^3 (a b)^4. \end{aligned}$$

Note that if $y \leq 1$, $B_{\xi}(x, y)$ and $B_{\xi'}(x, y)$ have at most one b ; and if $x \leq 1$, then each of these blocks has at most one a .

We first determine ways that for $l > 6$, $ba^l b$ may occur in $B_\xi(x, y)$, for any (x, y) . Because of the preceding observation, it suffices to consider only $x, y \geq 2$. If $x + y \geq 6$, $B_\xi(x, y)$ can be decomposed into a concatenation of F_j, G_k , and H with $j, k \geq 4$ as follows. Since $\xi(x, y) = 0$ for all $x + y \geq 5$, $B_\xi(x, y) = B_\xi(x, y - 1)B_\xi(x - 1, y)$. Continue factoring in this manner, reducing x and y , until each block in the factorization is one of $B_\xi(3, 3) = H, B_\xi(j, 2) = F_j$, or $B_\xi(2, k) = G_k$, for some $j, k \geq 4$. The concatenation that we arrive at must have the following structure. F_4 can be followed only by H . For $j > 4$, F_j can be followed only by F_{j-1} . G_4 can be preceded only by H . For $k > 4$, G_k can be preceded only by G_{k-1} . Note that for all $k, j \geq 4$ the concatenation $G_k F_j$ does occur in some $B_\xi(x, y)$, for example in $B_\xi(j + 1, k + 1)$.

None of the blocks H, F_j, G_k end with a^2 ; for all $l > 2, k \geq 4$, G_k and H do not contain a^l . So if $l > 6$ and $ba^l b$ appears in $B_\xi(x, y)$, then the majority of the a^l portion of $ba^l b$ must appear within one of the factors F_j of $B_\xi(x, y)$ for some $j \geq 4$. We can then see that $ba^l b$ appears in one of only four possible ways:

- (1) across a concatenation $G_k F_{l+1}$ for some $k \geq 4$,
- (2) across a concatenation $H F_{l+1}$,
- (3) across a concatenation $F_{l+1} F_l$,
- (4) entirely within a block F_j for some $j > l$.

In the first two cases, the appearance of $ba^l b$ is within the block $ba^l ba^l ba^{l-1} b$, while in the second two cases the appearance of $ba^l b$ is within the block $ba^l ba^{l-1} b$. Therefore, $ba^l b$ appears in any sequence in Σ_ξ only as a subblock of one of the two longer blocks $ba^l ba^l ba^{l-1} b$ or $ba^l ba^{l-1} b$.

We next determine ways that for $l > 6$, $ba^l b$ may occur in $B_{\xi'}(x, y)$, for any (x, y) . Again, it suffices to consider only $x, y \geq 2$. If $x + y \geq 6$, $B_{\xi'}(x, y)$ can be decomposed into a concatenation of F'_j, G'_k and H' with $j, k \geq 4$ as follows. Since $\xi'(x, y) = 1$ for all $x + y \geq 5$, $B_{\xi'}(x, y) = B_{\xi'}(x - 1, y)B_{\xi'}(x, y - 1)$. Continue factoring in this manner, reducing x and y , until each block in the factorization is one of $B'_{\xi}(3, 3) = H', B'_{\xi}(j, 2) = F'_j$, or $B'_{\xi}(2, k) = G'_k$, for some $j, k \geq 4$. The concatenation that we arrive at must have the following structure. F'_4 can be preceded only by H' . For $j > 4$, F'_j can be preceded only by F'_{j-1} . G'_4 can be followed only by H' . For $k > 4$, G'_k can be followed only by G'_{k-1} . Note that for all $k, j \geq 4$ the concatenation $F'_j G'_k$ does occur in some $B_{\xi'}(x, y)$, for example, in $B_{\xi'}(k + 1, j + 1)$.

None of the blocks H', F'_j, G'_k begin with a^2 ; for all $l > 2, k \geq 4$, G'_k and H' do not contain a^l . So if $l > 6$ and $ba^l b$ appears in $B_{\xi'}(x, y)$, then the majority of the a^l portion of $ba^l b$ must appear within one of the factors F'_j of $B_{\xi'}(x, y)$ for some $j \geq 4$. We can then see that $ba^l b$ appears in one of only four possible ways:

- (1) across a concatenation $F'_{l+1}G'_k$ for some $k \geq 4$,
- (2) across a concatenation $F'_{l+1}H$,
- (3) across a concatenation $F'_lF'_{l+1}$,
- (4) entirely within a block F'_j for some $j > l$.

In the first case, the appearance of $ba^l b$ is within the block $ba^l ba^l b^{k-1} a$ for some $k \geq 4$, while in the second case we see that the appearance of $ba^l b$ is within the block $ba^l ba^l b^2 ab$. In the third case the appearance of $ba^l b$ is within the block $ba^{l-1} ba^l bab$. In the final case if $j > l+1$ the appearance of $ba^l b$ is within the block $ba^{l-1} ba^l ba^{l+1} b$. However, if $j = l+1$, F'_{l+1} is not long enough to gather enough information about how $ba^l b$ is extended to the right and we must examine which blocks can follow F'_{l+1} . These are G'_k, H' , and F'_{l+2} . These give rise to the longer words $ba^l ba^l b^{k-1} a$ for some $k \geq 4$, $ba^l ba^l b^2 a$, and $ba^{l-1} ba^l ba^{l+1} b$, respectively. Therefore, $ba^l b$ appears in any sequence in $\Sigma_{\xi'}$ only as a subblock of one of these four longer words. Most significantly, we see that any extension of $ba^l b$ in $\mathcal{L}(\Sigma_{\xi'})$ will not be in $\mathcal{L}(\Sigma_{\xi'})$.

We use a similar argument to determine ways that, for $l > 6$, $ab^l a$ may occur in any $B_{\xi}(x, y)$ with $x, y \geq 2$. We find that $ab^l a$ appears in any sequence in Σ_{ξ} only at the end of the longer block $ab^{l-1} ab^l ab^l a$ or as a subblock of $ab^{l-1} ab^l a$.

Similarly, for $l > 6$, $ab^l a$ may occur in $B_{\xi'}(x, y)$ for $x, y \geq 2$ only as a subblock of one of the four longer blocks $ba^{j-1} b^l ab^l ab^{l-1}$ for some $j \geq 4$, $ba^2 b^l ab^l ab^{l-1} a$, $babab^l ab^{l-1} a$, or $ab^{l+1} ab^l ab^{l-1} a$. Most significantly, we see that any extension of $ab^l a$ in $\mathcal{L}(\Sigma_{\xi'})$ will not be in $\mathcal{L}(\Sigma_{\xi'})$.

Let $\omega \in \Sigma_{\xi} \cap \Sigma_{\xi'}$ and suppose that ω is not in the orbit of any of the sequences

$$a^{\infty}, b^{\infty}, a^{\infty}.ba^{\infty}, b^{\infty}.ab^{\infty}.$$

We will show that then a block $ba^l b$ or $ab^l a$ with $l > 6$ appears in ω , contradicting what was just established above.

Since $\omega = \dots \omega_{-2} \omega_{-1} \omega_0 \omega_1 \dots$ is not in one of the four special orbits listed above, there are at least three distinct values of i for which $\omega_i \neq \omega_{i+1}$, say for $i_1 < i_2 < i_3$; in other words, ω contains at least 3 “switches” from a to b or vice versa. Let $S = \omega_{i_1} \dots \omega_{i_3+1}$ be such a block which contains exactly three switches, so that S has the form $ba^m b^n a$ or $ab^n a^m b$ for some $m, n \geq 1$. Let $D = \omega_{i_1-100} \dots \omega_{i_1} \dots \omega_{i_3+101}$; in other words, D extends in ω to the left and the right of the three places i_1, i_2, i_3 , where switches are known to occur, by 100 entries. In addition, let E be the subblock of ω that extends D by 20 letters to the left and 20 letters to the right. Since $E \in \mathcal{L}(\Sigma_{\xi})$, there is a block $B_{\xi}(x, y)$ that contains E (with $x, y \geq 2$, since otherwise $B_{\xi}(x, y)$ contains only two switches). Moreover, the block H does not appear in ω , so E does not contain the entirety of H as a subblock. Therefore,

when $B_\xi(x, y)$ is written according to the above scheme as a concatenation of H, F_j, G_k for certain choices of $j, k \geq 4$, the portion of $B_\xi(x, y)$ containing D does not overlap any part of an appearance of H in the factorization (meaning that the intervals of indices of $B_\xi(x, y)$ at which any copies of D and H appear are disjoint). In other words, D is contained in its entirety within concatenations of the blocks F_j and G_k according to the structure given above.

Consider the factorization of $B_\xi(x, y)$ as described above, with $j, k \geq 4$. The block $S = \omega_{i_1} \dots \omega_{i_3+1}$ must overlap some blocks F_j and G_k of the factorization, and this can happen only with $j, k \geq 8$. This is because F_7 can appear in any sequence in Σ_ξ only as part of the string $F_7F_6F_5F_4H$; but we know that D does not overlap any portion of H , and the length of $F_7F_6F_5F_4$ is 100. Similarly, S cannot overlap F_6, F_5 , or F_4 . And if S overlaps a block G_k of the factorization, it must happen with $k \geq 8$, since G_7 can occur in any sequence in Σ_ξ only at the end of a string $HG_4G_5G_6G_7$, which also has length 100; and similarly for G_6, G_5 , and G_4 .

As mentioned above, S is either of the form ba^mb^na or ab^na^mb . If either m or n is greater than 6, then ba^lb or ab^la , with $l > 6$, appears in S , a subblock of D , a subblock of ω .

Suppose that both $m, n < 7$ and S overlaps F_j for some $j \geq 8$ or S overlaps G_k for some $k \geq 8$. Then inspection of the blocks F_j and G_k with $j, k \geq 8$ shows that S must necessarily overlap the suffix of F_j of length 45, or the prefix of length 45 of G_k , or both, for some $j, k \geq 8$. However, if S overlaps the last 45 entries of an appearance of a block F_j with $j \geq 8$, then the block ba^7b appears in the block $\omega_{i_1-45} \dots \omega_{i_1}$. Similarly, if S overlaps the first 45 entries of an appearance of a block G_k with $k \geq 8$, then the block ab^7a appears in the block $\omega_{i_3+1} \dots \omega_{i_3+45}$. Thus at least one of the blocks ba^7b or ab^7a appears in ω . We conclude that it is impossible that any $\omega \in \Sigma_\xi \cap \Sigma_{\xi'}$ is not in the orbit of one of the sequences $a^\infty, b^\infty, a^\infty.ba^\infty, b^\infty.ab^\infty$. \square

7. REMARKS, EXAMPLES, QUESTIONS

Question 7.1. Is coding orbits by the *first* edge also essentially faithful for every ordering of the Pascal diagram? In other words, under the hypotheses of Theorem 3.6, is the coding of γ by initial segments of length 1 not the same as the coding of γ' by initial segments of length 1?

Example 7.2. It is possible to assign an ordering to the Pascal graph in such a way that there are uncountably many minimal paths. It suffices to find a binary tree embedded in the graph, for then we can define the ordering in such a way that all edges in the tree are minimal, the other edges to vertices in the tree are maximal, and the orderings at

remaining vertices are assigned arbitrarily. Let the tree include the vertex $(0, 0)$ at level 0 and nonintersecting paths to vertices $(3, 0)$ and $(1, 2)$ at level 3.

Continue inductively as follows. Suppose that $n \geq 2$ and the tree from level 0 to level $2^n - 1$ contains every other vertex at level $2^n - 1$ and no other vertices at that level or at higher levels. (If these vertices are labeled $v_0 = (2^n - 1, 0), v_1 = (2^n - 2, 1), \dots, v_{2^n - 1} = (0, 2^n - 1)$, we are including $v_0, v_2, \dots, v_{2^n - 2}$.) Extend the tree to contain 2^n nonintersecting paths from these vertices to every fourth vertex at level $2^{n+1} - 2$. Then from each of these vertices at level $2^{n+1} - 2$ extend two paths to form nonintersecting paths to every other vertex at level 2^{n+1} .

We can simultaneously embed another dyadic tree which shares no edges with the first tree—if necessary, spread out the construction of the first tree even more. Then declaring that all edges in the second tree are maximal will produce an ordering for which there are uncountably many maximal paths as well as uncountably many minimal paths.

Question 7.3. We do not know whether it is possible for an ordering to have countably many maximal paths and uncountably many minimal paths, or vice versa. (But no ordering on the Pascal graph can have only finitely many maximal or minimal paths, since from each vertex there is a unique path up to the root consisting entirely of maximal (or entirely of minimal) edges.)

Question 7.4. We are interested in asymptotic estimates of the complexity function $p_\xi(n)$ which counts the number of n -blocks seen in the 1-codings of all sequences in the adic system (X, T_ξ) on the Pascal graph with a fixed ordering ξ . It is known [23] that for the ordinary Pascal system, determined by $\xi \equiv 0$, $p_0(n) \sim n^3/6$. Can some orderings produce asymptotically larger or smaller complexity functions? If orderings are chosen at random, for example by independent choices at each vertex of 0 with some probability $q \in (0, 1)$ and 1 with probability $1 - q$, what is the expected asymptotic complexity of the set of resulting codings?

Question 7.5. Our proofs of Theorem 3.6 and Theorem 4.2 depend heavily on particular properties of the Pascal graph. Exactly which properties are needed for these arguments to work? For some important dynamical properties (essential expansiveness, topological weak mixing, ...) could there be corresponding classes of Bratteli diagrams for which the properties hold for all adic transformations determined by all orderings?

Acknowledgements. We thank the referee for a careful and thorough reading and many helpful comments, corrections, and questions.

REFERENCES

- [1] Terrence M. Adams and Karl E. Petersen, *Binomial-coefficient multiples of irrationals*, *Monatsh. Math.* **125** (1998), no. 4, 269–278, DOI 10.1007/BF01305342. MR1621682 (99d:28026)
- [2] S. Bezuglyi, J. Kwiatkowski, and K. Medynets, *Aperiodic substitution systems and their Bratteli diagrams*, *Ergodic Theory Dynam. Systems* **29** (2009), no. 1, 37–72, DOI 10.1017/S0143385708000230. MR2470626 (2009m:37020)
- [3] S. Bezuglyi, J. Kwiatkowski, K. Medynets, and B. Solomyak, *Invariant measures on stationary Bratteli diagrams*, *Ergodic Theory Dynam. Systems* **30** (2010), no. 4, 973–1007, DOI 10.1017/S0143385709000443. MR2669408 (2012g:37019)
- [4] S. Bezuglyi, J. Kwiatkowski, and R. Yassawi, *Perfect orderings on finite rank Bratteli diagrams*, *Canad. J. Math.* **66** (2014), no. 1, 57–101, DOI 10.4153/CJM-2013-041-6. MR3150704
- [5] Sergey Bezuglyi and Reem Yassawi, *Perfect orderings on Bratteli diagrams II: general Bratteli diagrams*, arXiv:1306.1788 **342** (August 8, 2014) (English).
- [6] Tomasz Downarowicz and Alejandro Maass, *Finite-rank Bratteli-Vershik diagrams are expansive*, *Ergodic Theory Dynam. Systems* **28** (2008), no. 3, 739–747, DOI 10.1017/S0143385707000673. MR2422014 (2009c:37007)
- [7] Fabien Durand, *Combinatorics on Bratteli diagrams and dynamical systems*, *Combinatorics, Automata and Number Theory*, *Encyclopedia Math. Appl.*, vol. 135, Cambridge Univ. Press, Cambridge, 2010, pp. 324–372. MR2759109 (2012h:37023)
- [8] F. Durand, B. Host, and C. Skau, *Substitutional dynamical systems, Bratteli diagrams and dimension groups*, *Ergodic Theory Dynam. Systems* **19** (1999), no. 4, 953–993, DOI 10.1017/S0143385799133947. MR1709427 (2000i:46062)
- [9] Fabien Durand and Reem Yassawi, 2014. personal communication.
- [10] A. H. Forrest, *K-groups associated with substitution minimal systems*, *Israel J. Math.* **98** (1997), 101–139, DOI 10.1007/BF02937330. MR1459849 (99c:54056)
- [11] Sarah Bailey Frick, *Limited scope adic transformations*, *Discrete Contin. Dyn. Syst. Ser. S* **2** (2009), no. 2, 269–285, DOI 10.3934/dcdss.2009.2.269. MR2505638 (2010h:37001)
- [12] Thierry Giordano, Ian F. Putnam, and Christian F. Skau, *Topological orbit equivalence and C^* -crossed products*, *J. Reine Angew. Math.* **469** (1995), 51–111. MR1363826 (97g:46085)
- [13] Richard Gjerde and Ørjan Johansen, *Bratteli-Vershik models for Cantor minimal systems: applications to Toeplitz flows*, *Ergodic Theory Dynam. Systems* **20** (2000), no. 6, 1687–1710, DOI 10.1017/S0143385700000948. MR1804953 (2002a:46092)
- [14] F. J. Hahn, *Skew product transformations and the algebras generated by $\exp(p(n))$* , *Illinois J. Math.* **9** (1965), 178–190. MR0174989 (30 #5177)
- [15] Richard H. Herman, Ian F. Putnam, and Christian F. Skau, *Ordered Bratteli diagrams, dimension groups and topological dynamics*, *Internat. J. Math.* **3** (1992), no. 6, 827–864, DOI 10.1142/S0129167X92000382. MR1194074 (94f:46096)
- [16] Bernard Host, *Substitution subshifts and Bratteli diagrams*, *Topics in Symbolic Dynamics and Applications* (Temuco, 1997), *London Math. Soc. Lecture Note Ser.*, vol. 279, Cambridge Univ. Press, Cambridge, 2000, pp. 35–55. MR1776755 (2001e:37014)
- [17] Siri-Malén Høyne, *Finite-rank Bratteli-Vershik systems are expansive—a new proof*, arXiv:1411.3371 (November 12, 2014) (English).
- [18] Jeanette Janssen, Anthony Quas, and Reem Yassawi, *Bratteli diagrams where almost all orders are imperfect*, arXiv:1407.3496v2 (October 23, 2014) (English).
- [19] Harvey B. Keynes and James B. Robertson, *Eigenvalue theorems in topological transformation groups*, *Trans. Amer. Math. Soc.* **139** (1969), 359–369. MR0237748 (38 #6029)

- [20] A. Livshitz, *A sufficient condition for weak mixing of substitutions and stationary adic transformations*, Math. Notes **44** (1988), 920–925 (English, Russian).
- [21] Konstantin Medynets, *Cantor aperiodic systems and Bratteli diagrams*, C. R. Math. Acad. Sci. Paris **342** (2006), no. 1, 43–46, DOI 10.1016/j.crma.2005.10.024 (English, with English and French summaries). MR2193394 (2006g:37011)
- [22] Xavier Sebastien Méla, *Dynamical properties of the Pascal adic and related systems*, ProQuest LLC, Ann Arbor, MI, 2002. Thesis (Ph.D.)—The University of North Carolina at Chapel Hill. MR2704022
- [23] Xavier Méla and Karl Petersen, *Dynamical properties of the Pascal adic transformation*, Ergodic Theory Dynam. Systems **25** (2005), no. 1, 227–256, DOI 10.1017/S0143385704000173. MR2122921 (2005k:37012)

DEPARTMENT OF MATHEMATICS, FURMAN UNIVERSITY, GREENVILLE, SC 29613 USA

E-mail address: sarah.frick@furman.edu

DEPARTMENT OF MATHEMATICS, CB 3250 PHILLIPS HALL, UNIVERSITY OF NORTH CAROLINA, CHAPEL HILL, NC 27599 USA

E-mail address: petersen@math.unc.edu

COLLEGE OF CHARLESTON, 66 GEORGE ST., CHARLESTON, SC 29424-0001 USA

E-mail address: shieldss@cofc.edu